

Datasheet: MCA1940GA

Description:	MOUSE ANTI HUMAN CD19
Specificity:	CD19
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	LT19
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG from ascites prepared by precipitation and ion exchange chromatography.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	<p>UniProt: P15391 Related reagents</p> <p>Entrez Gene: 930 CD19 Related reagents</p>

Specificity **Mouse anti Human CD19 antibody, clone LT19** recognizes human CD19 also known as T-cell surface antigen Leu-12 or B-lymphocyte surface antigen B4. CD19 is a ~95 kDa type I single pass transmembrane glycoprotein expressed on follicular dendritic cells and B-cells during maturation but is lost on development into plasma cells ([de Rie et al. 1989](#)).

CD19 is the broadest lineage specific marker for B cells and functions as a B-cell co-receptor in conjunction with CD21 ([Bradbury et al. 1992](#)), CD9, CD81 and CD82 ([Horváth et al. 1998](#)). CD19 is implicated in the down-regulation of B cell growth and proliferation ([Pezzutto et al. 1987](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Hughes, G.J. *et al.* (2007) Virus immunocapture provides evidence of CD8 lymphocyte-derived HIV-1 *in vivo*. [AIDS. 21: 1507-13.](#)
2. Allen, J.S. *et al.* (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. [Diabetes. 58: 138-45.](#)
3. McIntosh, K. *et al.* (2006) The immunogenicity of human adipose-derived cells: temporal changes *in vitro*. [Stem Cells. 24: 1246-53.](#)
4. Sengstake, S. *et al.* (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. [Int Immunol. 18 \(7\): 1171-8.](#)
5. Villarroel Dorrego, M. *et al.* (2006) Transfection of CD40 in a human oral squamous cell carcinoma keratinocyte line upregulates immune potency and costimulatory molecules. [Br J Dermatol. 154: 231-8.](#)
6. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. [Blood. 118: 348-57.](#)
7. Lacal, P.M. *et al.* (2013) Glucocorticoid-induced tumor necrosis factor receptor family-related ligand triggering upregulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 and promotes leukocyte adhesion. [J Pharmacol Exp Ther. 347: 164-72.](#)
8. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. [Blood. 118: 348-57.](#)
9. Girbl, T. *et al.* (2013) CD40-mediated activation of chronic lymphocytic leukemia cells promotes their CD44-dependent adhesion to hyaluronan and restricts CCL21-induced motility. [Cancer Res. 73: 561-70.](#)
10. Hertzberg, L. *et al.* (2010) Down syndrome acute lymphoblastic leukemia, a highly heterogeneous disease in which aberrant expression of CRLF2 is associated with mutated JAK2: a report from the International BFM Study Group. [Blood. 115: 1006-17.](#)
11. Kakko, T. *et al.* (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. [Diab Vasc Dis Res. 8: 221-8.](#)
12. Dorvignit, D. *et al.* (2012) Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: a case study. [MAbs. 4 \(4\): 488-96.](#)
13. Karlsen, M. *et al.* (2015) TLR-7 and -9 Stimulation of Peripheral Blood B Cells Indicate Altered TLR Signalling in Primary Sjögren's Syndrome Patients by Increased Secretion of Cytokines. [Scand J Immunol. 82 \(6\): 523-31.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@549](#),
[DyLight@649](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Human Anti Mouse IgG1 (HCA036...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M288778:160607'

Printed on 02 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)