

Datasheet: MCA1960GA

Description:	MOUSE ANTI HUMAN CD200
Specificity:	CD200
Other names:	OX2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OX-104
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	UniProt: P41217 Related reagents

Entrez Gene:[4345](#) CD200 [Related reagents](#)

Synonyms MOX1, MOX2

Specificity **Mouse anti Human CD200 antibody, clone OX-104** recognizes the human CD200 cell surface antigen, also known as OX2.

CD200 is expressed by a subset of B lymphocytes, some endothelial cells and by neurons. Studies have suggested that the CD200-CD200 ligand system is of importance in the control of macrophage and granulocyte activation.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Wright, G.J. *et al.* (2001) The unusual distribution of the neuronal/lymphoid cell surface CD200 (OX2) glycoprotein is conserved in humans. [Immunology 102 \(2\): 173-9.](#)
2. Ko, Y.C. *et al.* (2009) Endothelial CD200 is heterogeneously distributed, regulated and involved in immune cell-endothelium interactions. [J Anat. 214: 183-95.](#)
3. Koning, N. *et al.* (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. [J Neuropathol Exp Neurol. 68: 159-67.](#)
4. Koning, N. *et al.* (2007) Downregulation of macrophage inhibitory molecules in multiple sclerosis lesions. [Ann Neurol. 62: 504-14.](#)
5. Raftery, M.J. *et al.* (2004) Shaping phenotype, function, and survival of dendritic cells by cytomegalovirus-encoded IL-10. [J Immunol. 173: 3383-91.](#)
6. Meuth, S.G. *et al.* (2008) CNS inflammation and neuronal degeneration is aggravated by impaired CD200-CD200R-mediated macrophage silencing. [J Neuroimmunol. 194: 62-9.](#)
7. Yamauchi, K. and Kurosaka, A. (2010) Expression and function of glycogen synthase kinase-3 in human hair follicles. [Arch Dermatol Res. 302: 263-70.](#)
8. Kloepper, J.E. *et al.* (2008) Immunophenotyping of the human bulge region: the quest to define useful *in situ* markers for human epithelial hair follicle stem cells and their niche. [Exp Dermatol. 17: 592-609.](#)
9. Ohyama, M. *et al.* (2006) Characterization and isolation of stem cell-enriched human hair follicle bulge cells. [J Clin Invest. 116: 249-60.](#)
10. Darmochwal-Kolarz, D. *et al.* (2013) The expressions of co-stimulatory molecules are altered on putative antigen-presenting cells in cord blood. [Am J Reprod Immunol. 69 \(2\): 180-7.](#)
11. Colmont, C.S. *et al.* (2013) CD200-expressing human basal cell carcinoma cells initiate tumor growth. [Proc Natl Acad Sci U S A. 110 \(4\): 1434-9.](#)
12. Chen, H.J. *et al.* (2015) Human placenta-derived adherent cells improve cardiac performance in mice with chronic heart failure. [Stem Cells Transl Med. 4 \(3\): 269-75.](#)
13. Ohyama, M. & Kobayashi, T. (2012) Isolation and characterization of stem cell-enriched human and canine hair follicle keratinocytes. [Methods Mol Biol. 879: 389-401.](#)
14. Patel, G.K. *et al.* (2012) Identification and characterization of tumor-initiating cells in human primary cutaneous squamous cell carcinoma. [J Invest Dermatol. 132 \(2\): 401-9.](#)
15. Kloepper, J.E. *et al.* (2008) Immunophenotyping of the human bulge region: the quest to define useful *in situ* markers for human epithelial hair follicle stem cells and their niche. [Exp Dermatol. 17 \(7\): 592-609.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@549 , DyLight@649 , DyLight@680 , DyLight@800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M319085:180719'

Printed on 01 Aug 2018