

## Datasheet: MCA1958F

<b>Description:</b>	RAT ANTI MOUSE CD200:FITC
<b>Specificity:</b>	CD200
<b>Other names:</b>	OX2
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-90
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
<b>Buffer Solution</b>	Phosphate buffered saline						
<b>Preservative</b>	0.09% Sodium Azide						
<b>Stabilisers</b>	1% Bovine Serum Albumin						
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1mg/ml						
<b>Immunogen</b>	Mouse CD200-rat CD4 fusion protein.						
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">O54901</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b>						

---

<b>Synonyms</b>	Mox2
-----------------	------

---

<b>Fusion Partners</b>	Spleen cells from immunised rats were fused with cells of the rat Y3 myeloma cell line
------------------------	--

---

<b>Specificity</b>	<p><b>Rat anti Mouse CD200 antibody, clone OX-90</b> recognizes the mouse CD200 cell surface antigen, also known as OX2.</p> <p>CD200 is expressed by splenic B lymphocytes, follicular dendritic cells, splenic endothelium and by neurons.</p> <p>Studies have suggested that the CD200 - CD200 ligand system is of importance in the control of macrophage and granulocyte activation.</p>
--------------------	---

---

<b>Flow Cytometry</b>	<p>Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<a href="#">BUF041A/B</a>).</p>
-----------------------	--

---

<b>References</b>	<ol style="list-style-type: none"><li>1. Hoek, R.M. <i>et al.</i> (2000) Down-regulation of the macrophage lineage through interaction with OX2 (CD200). <a href="#">Science. 290 (5497): 1768-71.</a></li><li>2. Nathan, C. &amp; Muller, W.A. (2001) Putting the brakes on innate immunity: a regulatory role for CD200? <a href="#">Nat Immunol. 2 (1): 17-9.</a></li><li>3. Broderick, C. <i>et al.</i> (2002) Constitutive retinal CD200 expression regulates resident microglia and activation state of inflammatory cells during experimental autoimmune uveoretinitis. <a href="#">Am J Pathol. 161 (5): 1669-77.</a></li><li>4. Seeds, R.E. <i>et al.</i> (2011) The role of myeloid receptors on murine plasmacytoid dendritic cells in induction of type I interferon. <a href="#">Int Immunopharmacol. 11 (7): 794-801.</a></li><li>5. Burger, P.E. <i>et al.</i> (2009) High aldehyde dehydrogenase activity: a novel functional marker of murine prostate stem/progenitor cells. <a href="#">Stem Cells. 27: 2220-8.</a></li><li>6. Garza, L.A. <i>et al.</i> (2011) Bald scalp in men with androgenetic alopecia retains hair follicle stem cells but lacks CD200-rich and CD34-positive hair follicle progenitor cells. <a href="#">J Clin Invest. 121: 613-22.</a></li><li>7. Ko, Y.C. <i>et al.</i> (2009) Endothelial CD200 is heterogeneously distributed, regulated and involved in immune cell-endothelium interactions. <a href="#">J Anat. 214: 183-95.</a></li><li>8. Koning, N. <i>et al.</i> (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. <a href="#">J Neuropathol Exp Neurol. 68: 159-67.</a></li><li>9. Krejsek, J. <i>et al.</i> (2010) Expression of CD200/CD200R regulatory molecules on granulocytes and monocytes is modulated by cardiac surgical operation. <a href="#">Perfusion. 25: 389-97.</a></li><li>10. Rijkers, E.S. (2007) Ligation of CD200R by CD200 is not required for normal murine myelopoiesis. <a href="#">Eur J Haematol. 79: 410-6.</a></li><li>11. Rijkers, E.S. <i>et al.</i> (2008) The inhibitory CD200R is differentially expressed on human and mouse T and B lymphocytes. <a href="#">Mol Immunol. 45: 1126-35.</a></li><li>12. Liu, J.Q. <i>et al.</i> (2016) A Critical Role for CD200R Signaling in Limiting the Growth and Metastasis of CD200+ Melanoma. <a href="#">J Immunol. Jul 6. pii: 1600052. [Epub ahead of print]</a></li><li>13. Montiel, M. <i>et al.</i> (2016) Melatonin decreases brain apoptosis, oxidative stress, and CD200 expression and increased survival rate in mice infected by Venezuelan equine encephalitis virus. <a href="#">Antivir Chem Chemother. Aug 8. pii: 2040206616660851. [Epub ahead of print]</a></li></ol>
-------------------	---

---

<b>Storage</b>	Store at +4°C or at -20°C if preferred.
----------------	---

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Shelf Life</b>	18 months from date of despatch.
-------------------	----------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: 10041: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M301406:170109'

**Printed on 01 Aug 2018**

---

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)