

Datasheet: MCA1957A700

Description:	RAT ANTI MOUSE CD68:Alexa Fluor® 700
Specificity:	CD68
Other names:	MACROSIALIN
Format:	ALEXA FLUOR® 700
Product Type:	Monoclonal Antibody
Clone:	FA-11
Isotype:	IgG2a
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	■			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) **Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (Product Code [BUF041](#)).

Target Species	Mouse		
Product Form	Purified IgG conjugated to Alexa Fluor® 700 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®700	702	723
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	Purified Concanavalin A acceptor glycoprotein from P815 cell line.		

UniProt:

[P31996](#) [Related reagents](#)

Entrez Gene:

[12514](#) Cd68 [Related reagents](#)

Specificity

Rat anti Mouse CD68 antibody, clone FA-11 recognizes mouse macrosialin, a heavily glycosylated transmembrane protein and murine homolog of human CD68, which is classified as a unique scavenger receptor (ScR) family member, due to the presence of a lysosome associated membrane protein (LAMP)-like domain.

CD68 is considered a pan macrophage marker, predominantly expressed on the intracellular lysosomes of tissue macrophages/monocytes, including Kupffer cells, microglia, histiocytes and osteoclasts, and is expressed to a lesser extent by dendritic cells and peripheral blood granulocytes.

CD68 is expressed by many tumor types including some B cell lymphomas, blastic NK lymphomas, melanomas, granulocytic (myeloid) sarcomas, hairy cell leukemias, and renal, urinary and pancreatic tumors, and can be used in cancer studies to demonstrate the presence/localization of macrophages.

Rat anti mouse CD68 antibody, clone FA-11, has been used in many mouse models for the identification of CD68 in immunohistochemical studies, using both frozen and paraffin-embedded tissues ([Masaki *et al.* 2003](#)) and ([Devey *et al.* 2009](#)).

Rat anti mouse CD68 antibody, clone FA-11 can be used in flow cytometry to detect intracellular CD68, following permeabilization, and can detect surface macrosialin at low levels in resident mouse peritoneal macrophages which can be enhanced with thioglycollate stimulation.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells in 100ul. Recommended protocols are available [Here](#)

References

1. Ramprasad, M.P. *et al.* (1996) Cell surface expression of mouse macrosialin and human CD68 and their role as macrophage receptors for oxidized low density lipoprotein. [Proc Natl Acad Sci U S A. 93 \(25\): 14833-8.](#)
2. Rabinowitz, S.S. & Gordon, S. (1991) Macrosialin, a macrophage-restricted membrane sialoprotein differentially glycosylated in response to inflammatory stimuli. [J Exp Med. 174 \(4\): 827-36.](#)
3. da Silva, R.P. & Gordon, S. (1999) Phagocytosis stimulates alternative glycosylation of macrosialin (mouse CD68), a macrophage-specific endosomal protein. [Biochem J. 338 \(Pt 3\): 687-94.](#)
4. Schleicher, U. *et al.* (2005) Minute numbers of contaminant CD8+ T cells or CD11b+CD11c+ NK cells are the source of IFN- γ in IL-12/IL-18-stimulated mouse macrophage populations. [Blood 105: 1319-1328.](#)
5. Choi, E.J. *et al.* (2014) Novel brain arteriovenous malformation mouse models for type 1 hereditary hemorrhagic telangiectasia. [PLoS One. 9\(2\): e88511.](#)
6. Kassim, S. *et al.* (2010) Gene therapy in a humanized mouse model of familial hypercholesterolemia leads to marked regression of atherosclerosis. [PloS ONE 5: e13424.](#)
7. Rahaman, S.O. *et al.* (2011) Vav family Rho guanine nucleotide exchange factors regulate CD36-mediated macrophage foam cell formation. [J Biol Chem. 286: 7010-7.](#)
8. Frossard, J.L. *et al.* (2011) Role of CCL-2, CCR-2 and CCR-4 in cerulein-induced acute pancreatitis and pancreatitis-associated lung injury. [J Clin Pathol. 64: 387-93](#)

9. West, E.L. *et al.* (2010) Long-term survival of photoreceptors transplanted into the adult murine neural retina requires immune modulation. [Stem Cells. 28: 1997-2007.](#)
10. Lopez, M.E. *et al.* (2011) Anatomically defined neuron-based rescue of neurodegenerative niemann-pick type C disorder. [J Neurosci. 31: 4367-78.](#)
11. Jayagopal, A. *et al.* (2009) Quantum dot mediated imaging of atherosclerosis. [Nanotechnology. 20: 165102.](#)
12. Leung, V.W. *et al.* (2009) Decay-accelerating factor suppresses complement C3 activation and retards atherosclerosis in low-density lipoprotein receptor-deficient mice. [Am J Pathol. 175: 1757-67.](#)
13. Devey, L. *et al.* (2009) Tissue-resident macrophages protect the liver from ischemia reperfusion injury via a heme oxygenase-1-dependent mechanism. [Mol Ther. 17: 65-72.](#)
14. Lu, W. *et al.* (2010) Photoacoustic imaging of living mouse brain vasculature using hollow gold nanospheres. [Biomaterials. 31: 2617-26.](#)
15. de Beer, M.C. *et al.* (2003) Lack of a direct role for macrosialin in oxidized LDL metabolism. [J Lipid Res. 44: 674-85.](#)
16. Song, L. *et al.* (2011) Deletion of the murine scavenger receptor CD68. [J Lipid Res. 52: 1542-50.](#)
17. Daldrup-Link, H.E. *et al.* (2011) MR Imaging of Tumor Associated Macrophages with Clinically-Applicable Iron Oxide Nanoparticles. [Clin Cancer Res. 17: 5695-704.](#)
18. Macauley, S.L. *et al.* (2011) The Role of Attenuated Astrocyte Activation in Infantile Neuronal Ceroid Lipofuscinosis [J. Neurosci 31: 15575-85.](#)
19. Martin-Manso, G. *et al.* (2008) Thrombospondin 1 promotes tumor macrophage recruitment and enhances tumor cell cytotoxicity of differentiated U937 cells. [Cancer Res. 68: 7090-9.](#)
20. Lazarini, F. *et al.* (2012) Early Activation of Microglia Triggers Long-Lasting Impairment of Adult Neurogenesis in the Olfactory Bulb [J Neurosci 32: 3652-64](#)
21. Hemmi, H. *et al.* (2009) A new triggering receptor expressed on myeloid cells (Trem) family member, Trem-like 4, binds to dead cells and is a DNAX activation protein 12-linked marker for subsets of mouse macrophages and dendritic cells. [J Immunol. 182: 1278-86.](#)
22. Akbarshahi, H. *et al.* (2012) Enrichment of Murine CD68(+)CCR2(+) and CD68(+)CD206(+) Lung Macrophages in Acute Pancreatitis-Associated Acute Lung Injury. [PLoS One. 7: e42654.](#)
23. Xiang, X. *et al.* (2016) TREM2 deficiency reduces the efficacy of immunotherapeutic amyloid clearance. [EMBO Mol Med. 8 \(9\): 992-1004.](#)
24. Masaki, T. *et al.* (2003) Heterogeneity of antigen expression explains controversy over glomerular macrophage accumulation in mouse glomerulonephritis. [Nephrol. Dial. Transplant 18:178-81.](#)
25. Dormishian, M. *et al.* (2013) Prokineticin receptor-1 is a new regulator of endothelial insulin uptake and capillary formation to control insulin sensitivity and cardiovascular and kidney functions. [J Am Heart Assoc. 2 \(5\): e000411.](#)
26. von Bargen, K. *et al.* (2014) Cervical Lymph Nodes as a Selective Niche for Brucella during Oral Infections. [PLoS One. 10 \(4\): e0121790.](#)
27. Hamour, S. *et al.* (2015) Local IL-17 Production Exerts a Protective Role in Murine Experimental Glomerulonephritis. [PLoS One. 10 \(8\): e0136238.](#)
28. Wang L *et al.* (2016) Bone Fracture Pre-Ischemic Stroke Exacerbates Ischemic Cerebral Injury in Mice. [PLoS One. 11 \(4\): e0153835.](#)
29. Nguyen, T.V. *et al.* (2016) Multiplex immunoassay characterization and species comparison of inflammation in acute and non-acute ischemic infarcts in human and mouse brain tissue. [Acta Neuropathol Commun. 4 \(1\): 100.](#)
30. Pena-Philippides, J.C. *et al.* (2016) *In vivo* inhibition of miR-155 significantly alters post-stroke inflammatory response. [J Neuroinflammation. 13 \(1\): 287.](#)
31. Paiva, A. A. *et al.* (2017) Apolipoprotein CIII Overexpression-Induced Hypertriglyceridemia Increases Nonalcoholic Fatty Liver Disease in Association with Inflammation and Cell Death. [Oxidative Med Cellular Longev. 2017: 1-18.](#)
32. Giraldo, J.A. *et al.* (2016) The impact of cell surface PEGylation and short-course

- immunotherapy on islet graft survival in an allogeneic murine model. [Acta Biomater. pii: S1742-7061\(16\)30656-0. \[Epub ahead of print\]](#)
33. Masuda, T. *et al.* (2017) Growth Factor Midkine Promotes Nuclear Factor of Activated T Cells-Regulated T-Cell-Activation and Th1 Cell Differentiation in Lupus Nephritis. [Am J Pathol. Feb 6. pii: S0002-9440\(17\)30029-9. \[Epub ahead of print\]](#)
34. Garofalo, S. *et al.* (2017) The glycoside oleandrin reduces glioma growth with direct and indirect effects on tumor cells. [J Neurosci. Mar 14. pii: 2296-16. \[Epub ahead of print\]](#)
35. Maeda, K. *et al.* (2017) Inhibition of H3K9 methyltransferase G9a ameliorates methylglyoxal-induced peritoneal fibrosis. [PLoS One. 12 \(3\): e0173706.](#)
36. Nishikawa, K. *et al.* (2015) Resveratrol increases CD68⁺ Kupffer cells colocalized with adipose differentiation-related protein and ameliorates high-fat-diet-induced fatty liver in mice. [Mol Nutr Food Res. 59 \(6\): 1155-70.](#)
37. Nagy, B. *et al.* (2017) Different patterns of neuronal activity trigger distinct responses of oligodendrocyte precursor cells in the corpus callosum. [PLoS Biol. 15 \(8\): e2001993.](#)
38. Koh, A.J. *et al.* (2017) The skeletal impact of the chemotherapeutic agent etoposide. [Osteoporos Int. 28 \(8\): 2321-33.](#)
39. Menzies, R.I. *et al.* (2017) Hyperglycemia-induced Renal P2X7 Receptor Activation Enhances Diabetes-related Injury. [EBioMedicine. 19: 73-83.](#)
40. Takane, K. *et al.* (2017) Detrimental Effects of Centrally Administered Angiotensin II are Enhanced in a Mouse Model of Alzheimer Disease Independently of Blood Pressure. [J Am Heart Assoc. 6 \(4\)Apr 20 \[Epub ahead of print\].](#)
41. Xuan, H. *et al.* (2017) Inhibition or deletion of angiotensin II type 1 receptor suppresses elastase-induced experimental abdominal aortic aneurysms. [J Vasc Surg. Apr 20 \[Epub ahead of print\].](#)
42. Metghalchi, S. *et al.* (2018) Indoleamine 2 3-dioxygenase knockout limits angiotensin II-induced aneurysm in low density lipoprotein receptor-deficient mice fed with high fat diet. [PLoS One. 13 \(3\): e0193737.](#)
43. Hill, N.R. *et al.* (2018) RIPK3-deficient mice were not protected from nephrotoxic nephritis. [BMC Nephrol. 19 \(1\): 61.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety

Material Safety Datasheet documentation #10041 available at:

Information 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:Alexa Fluor® 700 \(MCA1212A700\)](#)

Recommended Useful Reagents

[LEUCOPERM™ \(BUF09\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M299013:161215'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)