

Datasheet: MCA1957PE

Description:	RAT ANTI MOUSE CD68:RPE
Specificity:	CD68
Other names:	MACROSIALIN
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	FA-11
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	■			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) **Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (Product Code [BUF041](#)).

Target Species	Mouse		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1.0 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1%	Bovine Serum Albumin	
	5%	Sucrose	
Immunogen	Purified Concanavalin A acceptor glycoprotein from P815 cell line.		

UniProt:

[P31996](#) [Related reagents](#)

Entrez Gene:

[12514](#) Cd68 [Related reagents](#)

Specificity

Rat anti Mouse CD68 antibody, clone FA-11 recognizes mouse macrosialin, a heavily glycosylated transmembrane protein and murine homolog of human CD68, which is classified as a unique scavenger receptor (ScR) family member, due to the presence of a lysosome associated membrane protein (LAMP)-like domain.

CD68 is considered a pan macrophage marker, predominantly expressed on the intracellular lysosomes of tissue macrophages/monocytes, including Kupffer cells, microglia, histiocytes and osteoclasts, and is expressed to a lesser extent by dendritic cells and peripheral blood granulocytes.

CD68 is expressed by many tumor types including some B cell lymphomas, blastic NK lymphomas, melanomas, granulocytic (myeloid) sarcomas, hairy cell leukemias, and renal, urinary and pancreatic tumors, and can be used in cancer studies to demonstrate the presence/localization of macrophages.

Rat anti mouse CD68 antibody, clone FA-11, has been used in many mouse models for the identification of CD68 in immunohistochemical studies, using both frozen and paraffin-embedded tissues ([Masaki *et al.* 2003](#)) and ([Devey *et al.* 2009](#)).

Rat anti mouse CD68 antibody, clone FA-11 can be used in flow cytometry to detect intracellular CD68, following permeabilization, and can detect surface macrosialin at low levels in resident mouse peritoneal macrophages which can be enhanced with thioglycollate stimulation.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells in 100ul. Recommended protocols are available [Here](#)

References

1. Ramprasad, M.P. *et al.* (1996) Cell surface expression of mouse macrosialin and human CD68 and their role as macrophage receptors for oxidized low density lipoprotein. [Proc Natl Acad Sci U S A. 93 \(25\): 14833-8.](#)
2. Rabinowitz, S.S. & Gordon, S. (1991) Macrosialin, a macrophage-restricted membrane sialoprotein differentially glycosylated in response to inflammatory stimuli. [J Exp Med. 174 \(4\): 827-36.](#)
3. da Silva, R.P. & Gordon, S. (1999) Phagocytosis stimulates alternative glycosylation of macrosialin (mouse CD68), a macrophage-specific endosomal protein. [Biochem J. 338 \(Pt 3\): 687-94.](#)
4. Schleicher, U. *et al.* (2005) Minute numbers of contaminant CD8+ T cells or CD11b+CD11c+ NK cells are the source of IFN- γ in IL-12/IL-18-stimulated mouse macrophage populations. [Blood 105: 1319-1328.](#)
5. Choi, E.J. *et al.* (2014) Novel brain arteriovenous malformation mouse models for type 1 hereditary hemorrhagic telangiectasia. [PLoS One. 9\(2\): e88511.](#)
6. Kassim, S. *et al.* (2010) Gene therapy in a humanized mouse model of familial hypercholesterolemia leads to marked regression of atherosclerosis. [PloS ONE 5: e13424.](#)
7. Rahaman, S.O. *et al.* (2011) Vav family Rho guanine nucleotide exchange factors regulate CD36-mediated macrophage foam cell formation. [J Biol Chem. 286: 7010-7.](#)
8. Frossard, J.L. *et al.* (2011) Role of CCL-2, CCR-2 and CCR-4 in cerulein-induced acute pancreatitis and pancreatitis-associated lung injury. [J Clin Pathol. 64: 387-93](#)

9. West, E.L. *et al.* (2010) Long-term survival of photoreceptors transplanted into the adult murine neural retina requires immune modulation. [Stem Cells. 28: 1997-2007.](#)
10. Lopez, M.E. *et al.* (2011) Anatomically defined neuron-based rescue of neurodegenerative niemann-pick type C disorder. [J Neurosci. 31: 4367-78.](#)
11. Jayagopal, A. *et al.* (2009) Quantum dot mediated imaging of atherosclerosis. [Nanotechnology. 20: 165102.](#)
12. Leung, V.W. *et al.* (2009) Decay-accelerating factor suppresses complement C3 activation and retards atherosclerosis in low-density lipoprotein receptor-deficient mice. [Am J Pathol. 175: 1757-67.](#)
13. Devey, L. *et al.* (2009) Tissue-resident macrophages protect the liver from ischemia reperfusion injury via a heme oxygenase-1-dependent mechanism. [Mol Ther. 17: 65-72.](#)
14. Lu, W. *et al.* (2010) Photoacoustic imaging of living mouse brain vasculature using hollow gold nanospheres. [Biomaterials. 31: 2617-26.](#)
15. de Beer, M.C. *et al.* (2003) Lack of a direct role for macrosialin in oxidized LDL metabolism. [J Lipid Res. 44: 674-85.](#)
16. Song, L. *et al.* (2011) Deletion of the murine scavenger receptor CD68. [J Lipid Res. 52: 1542-50.](#)
17. Daldrup-Link, H.E. *et al.* (2011) MR Imaging of Tumor Associated Macrophages with Clinically-Applicable Iron Oxide Nanoparticles. [Clin Cancer Res. 17: 5695-704.](#)
18. Macauley, S.L. *et al.* (2011) The Role of Attenuated Astrocyte Activation in Infantile Neuronal Ceroid Lipofuscinosis [J. Neurosci 31: 15575-85.](#)
19. Martin-Manso, G. *et al.* (2008) Thrombospondin 1 promotes tumor macrophage recruitment and enhances tumor cell cytotoxicity of differentiated U937 cells. [Cancer Res. 68: 7090-9.](#)
20. Lazarini, F. *et al.* (2012) Early Activation of Microglia Triggers Long-Lasting Impairment of Adult Neurogenesis in the Olfactory Bulb [J Neurosci 32: 3652-64](#)
21. Hemmi, H. *et al.* (2009) A new triggering receptor expressed on myeloid cells (Trem) family member, Trem-like 4, binds to dead cells and is a DNAX activation protein 12-linked marker for subsets of mouse macrophages and dendritic cells. [J Immunol. 182: 1278-86.](#)
22. Akbarshahi, H. *et al.* (2012) Enrichment of Murine CD68(+)CCR2(+) and CD68(+)CD206(+) Lung Macrophages in Acute Pancreatitis-Associated Acute Lung Injury. [PLoS One. 7: e42654.](#)
23. Xiang, X. *et al.* (2016) TREM2 deficiency reduces the efficacy of immunotherapeutic amyloid clearance. [EMBO Mol Med. 8 \(9\): 992-1004.](#)
24. Masaki, T. *et al.* (2003) Heterogeneity of antigen expression explains controversy over glomerular macrophage accumulation in mouse glomerulonephritis. [Nephrol. Dial. Transplant 18:178-81.](#)
25. Dormishian, M. *et al.* (2013) Prokineticin receptor-1 is a new regulator of endothelial insulin uptake and capillary formation to control insulin sensitivity and cardiovascular and kidney functions. [J Am Heart Assoc. 2 \(5\): e000411.](#)
26. von Bargen, K. *et al.* (2014) Cervical Lymph Nodes as a Selective Niche for Brucella during Oral Infections. [PLoS One. 10 \(4\): e0121790.](#)
27. Hamour, S. *et al.* (2015) Local IL-17 Production Exerts a Protective Role in Murine Experimental Glomerulonephritis. [PLoS One. 10 \(8\): e0136238.](#)
28. Wang L *et al.* (2016) Bone Fracture Pre-Ischemic Stroke Exacerbates Ischemic Cerebral Injury in Mice. [PLoS One. 11 \(4\): e0153835.](#)
29. Nguyen, T.V. *et al.* (2016) Multiplex immunoassay characterization and species comparison of inflammation in acute and non-acute ischemic infarcts in human and mouse brain tissue. [Acta Neuropathol Commun. 4 \(1\): 100.](#)
30. Pena-Philippides, J.C. *et al.* (2016) *In vivo* inhibition of miR-155 significantly alters post-stroke inflammatory response. [J Neuroinflammation. 13 \(1\): 287.](#)
31. Paiva, A. A. *et al.* (2017) Apolipoprotein CIII Overexpression-Induced Hypertriglyceridemia Increases Nonalcoholic Fatty Liver Disease in Association with Inflammation and Cell Death. [Oxidative Med Cellular Longev. 2017: 1-18.](#)
32. Giraldo, J.A. *et al.* (2016) The impact of cell surface PEGylation and short-course

- immunotherapy on islet graft survival in an allogeneic murine model. [Acta Biomater. pii: S1742-7061\(16\)30656-0. \[Epub ahead of print\]](#)
33. Masuda, T. *et al.* (2017) Growth Factor Midkine Promotes Nuclear Factor of Activated T Cells-Regulated T-Cell-Activation and Th1 Cell Differentiation in Lupus Nephritis. [Am J Pathol. Feb 6. pii: S0002-9440\(17\)30029-9. \[Epub ahead of print\]](#)
34. Garofalo, S. *et al.* (2017) The glycoside oleandrin reduces glioma growth with direct and indirect effects on tumor cells. [J Neurosci. Mar 14. pii: 2296-16. \[Epub ahead of print\]](#)
35. Maeda, K. *et al.* (2017) Inhibition of H3K9 methyltransferase G9a ameliorates methylglyoxal-induced peritoneal fibrosis. [PLoS One. 12 \(3\): e0173706.](#)
36. Nishikawa, K. *et al.* (2015) Resveratrol increases CD68⁺ Kupffer cells colocalized with adipose differentiation-related protein and ameliorates high-fat-diet-induced fatty liver in mice. [Mol Nutr Food Res. 59 \(6\): 1155-70.](#)
37. Nagy, B. *et al.* (2017) Different patterns of neuronal activity trigger distinct responses of oligodendrocyte precursor cells in the corpus callosum. [PLoS Biol. 15 \(8\): e2001993.](#)
38. Koh, A.J. *et al.* (2017) The skeletal impact of the chemotherapeutic agent etoposide. [Osteoporos Int. 28 \(8\): 2321-33.](#)
39. Menzies, R.I. *et al.* (2017) Hyperglycemia-induced Renal P2X7 Receptor Activation Enhances Diabetes-related Injury. [EBioMedicine. 19: 73-83.](#)
40. Takane, K. *et al.* (2017) Detrimental Effects of Centrally Administered Angiotensin II are Enhanced in a Mouse Model of Alzheimer Disease Independently of Blood Pressure. [J Am Heart Assoc. 6 \(4\)Apr 20 \[Epub ahead of print\].](#)
41. Xuan, H. *et al.* (2017) Inhibition or deletion of angiotensin II type 1 receptor suppresses elastase-induced experimental abdominal aortic aneurysms. [J Vasc Surg. Apr 20 \[Epub ahead of print\].](#)
42. Metghalchi, S. *et al.* (2018) Indoleamine 2 3-dioxygenase knockout limits angiotensin II-induced aneurysm in low density lipoprotein receptor-deficient mice fed with high fat diet. [PLoS One. 13 \(3\): e0193737.](#)
43. Hill, N.R. *et al.* (2018) RIPK3-deficient mice were not protected from nephrotoxic nephritis. [BMC Nephrol. 19 \(1\): 61.](#)

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:RPE \(MCA1212PE\)](#)

Recommended Useful Reagents

[LEUCOPERM™ \(BUF09\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com
"M300980:170106"

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)