

Datasheet: MCA193F

Description:	MOUSE ANTI RAT IgE:FITC
Specificity:	IgE
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MARE-1
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			5ug/ml
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.1% Sodium Azide		
Stabilisers	50% Glycerol		
Approx. Protein Concentrations	IgG concentration 1 mg/ml		
Immunogen	Rat IR162, IR1016, IR2 and IR410 IgE myeloma proteins.		
External Database Links	UniProt:		
	P01855	Related reagents	

Entrez Gene:

[299351](#) Ighe [Related reagents](#)

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
------------------------	---

Specificity	<p>Mouse anti Rat IgE antibody, clone MARE-1 recognizes rat epsilon heavy chain of immunoglobulin IgE and does not cross-react with other classes of rat immunoglobulin.</p> <p>Mouse anti Rat IgE antibody, clone MARE-1 binds to rat IgE with an avidity of $4 \times 10^9 \text{M}^{-1}$</p>
--------------------	---

Flow Cytometry	Use 50ul of the suggested working dilution to label 2×10^7 cells in 100ul.
-----------------------	---

References	<ol style="list-style-type: none">1. Negrão-Corrêa, D. <i>et al.</i> (1996) Intestinal transport and catabolism of IgE: a major blood-independent pathway of IgE dissemination during a <i>Trichinella spiralis</i> infection of rats. J Immunol. 157 (9): 4037-44.2. Bazin, H. <i>et al.</i> (1984) Rat monoclonal antibodies. I. Rapid purification from <i>in vitro</i> culture supernatants. J Immunol Methods. 66 (2): 261-9.3. Bazin, H. <i>et al.</i> (1974) Transplantable immunoglobulin-secreting tumours in rats. IV. Sixty-three IgE-secreting immunocytoma tumours. Immunology. 26 (4): 713-23.4. Bazin, H. <i>et al.</i> (1978) Transplantable IgD immunoglobulin-secreting tumors in rat. J Immunol. 121 (5): 2077-82.5. Bazin, H. <i>et al.</i> (1984) Rat monoclonal antibodies. II. A rapid and efficient method of purification from ascitic fluid or serum. J Immunol Methods. 71 (1): 9-16.6. Cho, J.K. & Cho, S.W. (2000) Shared epitope for monoclonal IR162 between <i>Anisakis simplex</i> larvae and <i>Clonorchis sinensis</i> and cross-reactivity between antigens. J Parasitol. 86 (5): 1145-9.7. Silveira, M.R. <i>et al.</i> (2002) Infection with <i>Strongyloides venezuelensis</i> induces transient airway eosinophilic inflammation, an increase in immunoglobulin E, and hyperresponsiveness in rats. Infect Immun. 70: 6263-72.8. Korinek, M. <i>et al.</i> (2016) Anti-allergic potential of <i>Typhonium blumei</i>: inhibition of degranulation via suppression of PI3K/PLCγ2 phosphorylation and calcium influx. Phytomedicine. 23 (14): 1706-15.9. Bąbólewska, E. & Brzezińska-błaszczyk, E. (2015) Human-derived cathelicidin LL-37 directly activates mast cells to proinflammatory mediator synthesis and migratory response. Cell Immunol. 293 (2): 67-73.10. Agier, J. <i>et al.</i> (2018) Cathelicidin LL-37 Affects Surface and Intracellular Toll-Like Receptor Expression in Tissue Mast Cells. J Immunol Res. 2018: 7357162.
-------------------	--

Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10328 available at: 10328: https://www.bio-rad-antibodies.com/uploads/MSDS/10328.pdf
--------------------------------------	---

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA1209F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M314165:180412'

Printed on 16 May 2018