

Datasheet: MCA2059

Description:	MOUSE ANTI HUMAN CD88
Specificity:	CD88
Other names:	C5aR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	P12/1
Isotype:	IgG2a
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/50 - 1/100
Immunohistology - Frozen	■			
Immunohistology - Paraffin	■			
ELISA			■	
Immunoprecipitation		■		
Western Blotting	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Rhesus Monkey N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml

Immunogen	C5aR - peptide: Met ₁ - Asn ₃₁
External Database Links	UniProt: P21730 Related reagents Entrez Gene: 728 C5AR1 Related reagents
Synonyms	C5AR, C5R1
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63 - Ag8 myeloma cell line.
Specificity	Mouse anti Human CD88 antibody, clone P12/1 recognizes the C5a receptor (C5aR) also known as CD88 or C5a anaphylatoxin chemotactic receptor 1. CD88 is predominantly expressed on cells of the myeloid lineage. Mouse anti Human CD88 antibody, clone P12/1 does not inhibit the binding of C5a to its receptor.
Flow Cytometry	Use 10ul of the suggested working dilution to label 5 x 10 ⁵ cells in 100ul.
References	1. Oppermann, M. <i>et al.</i> (1993) Probing the human receptor for C5a anaphylatoxin with site-directed antibodies. Identification of a potential ligand binding site on the NH2-terminal domain. J Immunol. 151 (7): 3785-94. 2. Opperman, M. <i>et al.</i> (1995) Antibodies from the myeloid panel that react with the C5a receptor and antagonize C5a biological activity. In: Schlossman, S.F. (ed.) Leucocyte Typing V. OUP: 955-6. 3. Werfel, T. <i>et al.</i> (1995) Binding of anti-C5a receptor (C5aR) antibodies to cells of clinically normal human skin. In: Schlossman, S.F. (ed.) Leucocyte Typing V. OUP: 957-9. 4. Oppermann, M. (1995) Cluster report: (C5a receptor). In: Schlossman, S.F. (ed.) Leucocyte Typing V. OUP: 953-4. 5. Oppermann, M. & Götze, O. (1994) Plasma clearance of the human C5a anaphylatoxin by binding to leucocyte C5a receptors. Immunology. 82 (4): 516-21. 6. Werfel, T. <i>et al.</i> (1996) CD88 antibodies specifically bind to C5aR on dermal CD117+ and CD14+ cells and react with a desmosomal antigen in human skin. J Immunol. 157: 1729-35. 7. Morris, A.C. <i>et al.</i> (2011) C5a-mediated neutrophil dysfunction is RhoA-dependent and predicts infection in critically ill patients. Blood. 117: 5178-88. 8. Hüttenrauch, F. <i>et al.</i> (2005) G protein-coupled receptor kinases promote phosphorylation and beta-arrestin-mediated internalization of CCR5 homo- and hetero-oligomers. J Biol Chem. 280: 37503-15. 9. Nitta, H. <i>et al.</i> (2013) Enhancement of human cancer cell motility and invasiveness by anaphylatoxin C5a via aberrantly expressed C5a receptor (CD88). Clin Cancer Res. 19 (8): 2004-13. 10. Patzelt, J. <i>et al.</i> (2015) Expression of anaphylatoxin receptors on platelets in patients with coronary heart disease. Atherosclerosis. 238 (2): 289-95. 11. Huber-Lang, M. <i>et al.</i> (2005) Changes in the novel orphan, C5a receptor (C5L2), during experimental sepsis and sepsis in humans. J Immunol. 174 (2): 1104-10. 12. Unnewehr H <i>et al.</i> (2013) Changes and regulation of the C5a receptor on neutrophils during septic shock in humans. J Immunol. 190 (8): 4215-25. 13. Visser T <i>et al.</i> (2011) Isolated blunt chest injury leads to transient activation of circulating neutrophils. Eur J Trauma Emerg Surg. 37 (2): 177-184. 14. Visser T <i>et al.</i> (2012) Homology in systemic neutrophil response induced by human

- experimental endotoxemia and by trauma. [Shock. 37 \(2\): 145-51.](#)
15. Pollok-Kopp B *et al.* (2007) Dynamics of protein kinase C-mediated phosphorylation of the complement C5a receptor on serine 334. [J Biol Chem. 282 \(7\): 4345-53.](#)
16. Zannetti, C. *et al.* (2016) Characterization of the Inflammasome in Human Kupffer Cells in Response to Synthetic Agonists and Pathogens. [J Immunol. May 25. pii: 1502301. \[Epub ahead of print\]](#)
17. Presicce, P. *et al.* (2015) Neutrophil recruitment and activation in decidua with intra-amniotic IL-1beta in the preterm rhesus macaque. [Biol Reprod. 92 \(2\): 56.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Human Anti Mouse IgG2a (HCA037...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America

Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M315489:180503'

Printed on 18 May 2018

