

Datasheet: MCA2142APC

Description:	MOUSE ANTI HUMAN CD63:APC
Specificity:	CD63
Format:	APC
Product Type:	Monoclonal Antibody
Clone:	MEM-259
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Allophycocyanin (APC) - lyophilised		
Reconstitution	Reconstitute with 1ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	APC	650	661
Preparation	Purified IgG prepared by affinity chromatography on Protein A		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	HPB-ALL cell line.		
External Database Links	UniProt: P08962 Related reagents		

Entrez Gene:[967](#) CD63 [Related reagents](#)**Synonyms**

MLA1, TSPAN30

Specificity

Mouse anti Human CD63 antibody, clone MEM-259 recognizes the CD63 cell surface antigen, also known as granulophysin, lysosomal associated membrane protein 3 (LAMP-3), Melanoma-associated antigen ME491, Ocular melanoma-associated antigen or tetraspanin-30. CD63 is a 238 amino acid, tetraspanin glycoprotein of ~40-60 kDa.

CD63 is expressed by monocytes, macrophages, endothelial cells where it is a component of Weibel-Palade bodies ([Vischer et al. 1993](#)) and by a variety of cell lines. It is located intracellularly in lysosomal granules of platelets, being translocated to the surface upon activation. It may therefore be useful as a marker of platelet activation ([Hamamoto et al. 1994](#)). CD63 expression is also associated with melanoma development ([Radford et al. 1997](#)) and expression is rapidly down-regulated by progesterone in endometrial tissues ([Okada et al. 1999](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells in 100ul or 100ul whole blood.

References

1. Casey, T.M. *et al.* (2007) Organelle proteomics: identification of the exocytic machinery associated with the natural killer cell secretory lysosome. [Mol Cell Proteomics. 6 \(5\): 767-80.](#)
2. Durand-Panteix, S. *et al.* (2012) B7-H1, which represses EBV-immortalized B cell killing by autologous T and NK cells, is oppositely regulated by c-Myc and EBV latency III program at both mRNA and secretory lysosome levels. [J Immunol. 189 \(1\): 181-90.](#)
3. Abdel-Latif, D. *et al.* (2004) Rac2 is critical for neutrophil primary granule exocytosis. [Blood. 104: 832-9.](#)
4. Norling, L.V. *et al.* (2012) Resolvin D1 limits polymorphonuclear leukocyte recruitment to inflammatory loci: receptor-dependent actions. [Arterioscler Thromb Vasc Biol. 32 \(8\): 1970-8.](#)
5. McKechnie, N.M. *et al.* (2006) Fas-ligand is stored in secretory lysosomes of ocular barrier epithelia and released with microvesicles. [Exp Eye Res. 83: 304-14.](#)
6. Spring, F.A. *et al.* (2013) Tetraspanins CD81 and CD82 facilitate $\alpha\beta 1$ -mediated adhesion of human erythroblasts to vascular cell adhesion molecule-1. [PLoS One. 8\(5\):e62654.](#)
7. Pliyev, B.K. (2008) Activated human neutrophils rapidly release the chemotactically active D2D3 form of the urokinase-type plasminogen activator receptor (uPAR/CD87). [Mol Cell Biochem. 321: 111-22.](#)
8. Suraeva, N.M. *et al.* (2015) Changes in the Morphological and Immunological Characteristics of Mel Ibr Melanoma Cells in Response to Chicken Embryo Extract. [Bull Exp Biol Med. 159 \(4\): 520-3.](#)
9. Schäfer, T. *et al.* (2010) A granular variant of CD63 is a regulator of repeated human mast cell degranulation. [Allergy. 65 \(10\): 1242-55.](#)
10. Jakhria, T. *et al.* (2014) $\beta 2$ -microglobulin amyloid fibrils are nanoparticles that disrupt lysosomal membrane protein trafficking and inhibit protein degradation by lysosomes. [J Biol Chem. 289 \(52\): 35781-94.](#)
11. Ethier, C. *et al.* (2016) Calcitriol Reduces Eosinophil Necrosis Which Leads to the Diminished Release of Cytotoxic Granules. [Int Arch Allergy Immunol. 171 \(2\): 119-29.](#)

Storage

Prior to reconstitution store at +4°C.

After reconstitution store at +4°C.

DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:APC \(MCA928APC\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M316629:180529'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)