

## Datasheet: MCA2282A647

<b>Description:</b>	MOUSE ANTI HUMAN CD200R:Alexa Fluor® 647
<b>Specificity:</b>	CD200R
<b>Other names:</b>	OX2R
<b>Format:</b>	ALEXA FLUOR® 647
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX108
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/1ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human		
<b>Product Form</b>	Purified IgG conjugated to Alexa Fluor® 647 - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	Alexa Fluor®647	650	665
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative</b>	0.09% Sodium Azide		
<b>Stabilisers</b>	1% Bovine Serum Albumin		
<b>Approx. Protein Concentrations</b>	IgG concentration 0.05 mg/ml		
<b>Immunogen</b>	Fusion protein hCD200RCD4d3+4.		
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">Q8TD46</a> <a href="#">Related reagents</a>		

**Entrez Gene:**

[131450](#) CD200R1 [Related reagents](#)

---

**Synonyms** CD200R, CRTR2, MOX2R, OX2R

---

**Specificity** **Mouse anti Human CD200R antibody, clone OX108** recognizes human CD200R, a cell surface glycoprotein (also known as OX2R). In humans CD200R is expressed primarily by peripheral blood monocytes and neutrophils but also by other leucocytes including T-lymphocytes and mast cells, CD200-CD200R interaction may be involved in the control of myeloid cellular function ([Wright \*et al.\* 2003](#)).

Levels of expression on resting peripheral blood cells are relatively low.

---

**Flow Cytometry** Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul

- 
- References**
1. Wright, G.J. *et al.* (2003) Characterization of the CD200 receptor family in mice and humans and their interactions with CD200. [J Immunol. 171 \(6\): 3034-46.](#)
  2. Koning, N. *et al.* (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. [J Neuropathol Exp Neurol. 68: 159-67.](#)
  3. Melief, J. *et al.* (2013) Microglia in normal appearing white matter of multiple sclerosis are alerted but immunosuppressed. [Glia. 61 \(11\): 1848-61.](#)
  4. Ambarus, C.A. *et al.* (2012) Intimal lining layer macrophages but not synovial sublining macrophages display an IL-10 polarized-like phenotype in chronic synovitis. [Arthritis Res Ther. 14 \(2\): R74.](#)
  5. Holmannova, D. *et al.* (2015) Inhibitory CD200R and proapoptotic CD95/CD95L molecules on innate immunity cells are modulated by cardiac surgery. [Perfusion. 30 \(7\): 543-55.](#)
  6. Ambarus CA *et al.* (2012) Soluble immune complexes shift the TLR-induced cytokine production of distinct polarized human macrophage subsets towards IL-10. [PLoS One. 7 \(4\): e35994.](#)
  7. Memarian, A. *et al.* (2013) Upregulation of CD200 is associated with Foxp3+ regulatory T cell expansion and disease progression in acute myeloid leukemia. [Tumour Biol. 34 \(1\): 531-42.](#)
  8. Sousa, S. *et al.* (2015) Human breast cancer cells educate macrophages toward the M2 activation status. [Breast Cancer Res. 17 \(1\): 101.](#)
  9. Vicetti Miguel, R.D. *et al.* (2013) Human female genital tract infection by the obligate intracellular bacterium *Chlamydia trachomatis* elicits robust Type 2 immunity. [PLoS One. 8 \(3\): e58565.](#)
  10. van de Garde, M.D. *et al.* (2014) Chronic exposure to glucocorticoids shapes gene expression and modulates innate and adaptive activation pathways in macrophages with distinct changes in leukocyte attraction. [J Immunol. 192 \(3\): 1196-208.](#)
  11. Melief, J. *et al.* (2012) Phenotyping primary human microglia: tight regulation of LPS responsiveness. [Glia. 60 \(10\): 1506-17.](#)
  12. Georgoudaki, A.M. *et al.* (2016) Reprogramming Tumor-Associated Macrophages by Antibody Targeting Inhibits Cancer Progression and Metastasis. [Cell Rep. 15 \(9\): 2000-11.](#)
  13. BjörnfortHolmström, S. *et al.* (2017) Gingival Tissue Inflammation Promotes Increased Matrix Metalloproteinase-12 Production by CD200R<sup>low</sup> Monocyte-Derived Cells in Periodontitis. [J Immunol. Nov 03 \[Epub ahead of print\].](#)

---

**Storage** Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Shelf Life** 18 months from date of despatch.

---

**Acknowledgements** This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA928A647\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M316630:180529'

**Printed on 20 Jun 2018**