

Datasheet: MCA2297

Description:	RAT ANTI MOUSE CD106
Specificity:	CD106
Other names:	VCAM-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MVCAM A (429)
Isotype:	IgG2a
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/50
Immunohistology - Frozen	■			
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	
Immunofluorescence	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Stromal cell line PA6.

**External Database
Links**

UniProt:

[P29533](#) [Related reagents](#)

Entrez Gene:

[22329](#) Vcam1 [Related reagents](#)

Synonyms

Vcam-1

Fusion Partners

Spleen cells from immunised Lewis rats were fused with cells of the mouse P3X63Ag8.653 myeloma cell line.

Specificity

Rat anti Mouse CD106 antibody, clone MVCAM A (429) recognizes murine vascular adhesion molecule 1 (VCAM-1), a cell surface glycoprotein that is also known as CD106. CD106 is expressed predominantly on endothelial cells and expression is up-regulated during inflammation. The ligand for CD106 is the alpha 4 subunit (CD49d) of the integrin VLA-4 (CD49d/CD29).

Rat anti Mouse CD106 antibody, clone MVCAM A (429) is reported to partially block VCAM-1 mediated functions ([Kinashi and Springer 1994](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Kinashi, T. and Springer, T.A. (1994) Adhesion molecules in hematopoietic cells. [Blood Cells. 20:25-44.](#)
2. Wormley, F.L. Jr *et al.* (2001) Cell adhesion molecule and lymphocyte activation marker expression during experimental vaginal candidiasis. [Infect Immun. 69 \(8\): 5072-9.](#)
3. Woo, J.M. *et al.* (2010) Treatment with apolipoprotein A-1 mimetic peptide reduces lupus-like manifestations in a murine lupus model of accelerated atherosclerosis. [Arthritis Res Ther. 12\(3\):R93.](#)
4. Hall, L.J. *et al.* (2010) Probing local innate immune responses after mucosal immunisation. [J Immune Based Ther Vaccines. 8: 5.](#)
5. Winnik S *et al.* (2011) Dietary α-linolenic acid diminishes experimental atherogenesis and restricts T cell-driven inflammation. [Eur Heart J. 32 \(20\): 2573-84.](#)
6. Hamada, T. *et al.* (2009) Inducible nitric oxide synthase deficiency impairs matrix metalloproteinase-9 activity and disrupts leukocyte migration in hepatic ischemia/reperfusion injury. [Am J Pathol. 174: 2265-77.](#)
7. Hamada, T. *et al.* (2008) Metalloproteinase-9 deficiency protects against hepatic ischemia/reperfusion injury. [Hepatology. 47: 186-98.](#)
8. Baumer, Y. *et al.* (2011) Telomerase-based immortalization modifies the angiogenic/inflammatory responses of human coronary artery endothelial cells. [Exp Biol Med \(Maywood\). 236: 692-700.](#)
9. Lau, H.Y. and Bhatia, M. (2007) Effect of CP-96,345 on the expression of adhesion molecules in acute pancreatitis in mice. [Am J Physiol Gastrointest Liver Physiol. 292: G1283-92.](#)
10. Kumpers, P. *et al.* (2011) The synthetic tie2 agonist peptide vasculotide protects against vascular leakage and reduces mortality in murine abdominal sepsis. [Crit Care. 15 \(5\): R261.](#)
11. Püntener, U. *et al.* (2012) Long-term impact of systemic bacterial infection on the cerebral vasculature and microglia. [J Neuroinflammation. 9: 146.](#)
12. David, S. *et al.* (2011) Acute administration of recombinant Angiopoietin-1 ameliorates multiple-organ dysfunction syndrome and improves survival in murine sepsis. [Cytokine. 55 \(2\): 251-9.](#)
13. Kuriyama, N. *et al.* (2011) Tenascin-C: a novel mediator of hepatic ischemia and reperfusion injury. [Hepatology. 54 \(6\): 2125-36.](#)
14. Braach, N. *et al.* (2014) RAGE controls activation and anti-inflammatory signalling of protein C. [PLoS One. 9 \(2\): e89422.](#)

15. Li, M. *et al.* (2014) The indoleamine 2,3-dioxygenase pathway controls complement-dependent enhancement of chemo-radiation therapy against murine glioblastoma. [J Immunother Cancer. 2: 21.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®549 , DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M317140:180619'

Printed on 21 Jun 2018