

Datasheet: MCA2281GA

Description:	RAT ANTI MOUSE CD200R
Specificity:	CD200R
Other names:	OX2R
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OX-110
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/25 - 1/50
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Fusion protein mCD200RCD4d3+4

**External Database
Links**

UniProt:

[Q9ES57](#) [Related reagents](#)

Entrez Gene:

[57781](#) Cd200r1 [Related reagents](#)

Synonyms

Mox2r, Ox2r

Specificity

Rat anti Mouse CD200R antibody, clone OX-110 recognizes mouse CD200R, a cell surface glycoprotein (also known as OX2R. Mouse CD200R is a 326 amino acid, ~48 kDa single pass type I transmembrane glycoprotein, expressed primarily by peripheral blood monocytes and neutrophils but also by other leucocytes including T-lymphocytes and mast cells ([Wright et al.2003](#)). Studies suggest that CD200-CD200R interaction may be involved in the control of myeloid cellular function ([Minas and Liversidge 2006](#)).

Rat anti Mouse CD200R antibody, clone OX-110 has been used successfully for the immunohistochemical detection of CD200R expressing cells on acetone fixed cryosections of murine synovial tissue ([Simelyte et al. 2008](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul

References

1. Wright, G.J. *et al.* (2003) Characterization of the CD200 receptor family in mice and humans and their interactions with CD200. [J Immunol. 171 \(6\): 3034-46.](#)
2. Zhao, J. *et al.* (2009) Evasion by stealth: inefficient immune activation underlies poor T cell response and severe disease in SARS-CoV-infected mice. [PLoS Pathog. 5 \(10\): e1000636.](#)
3. Björck, P. *et al.* (2011) Plasmacytoid Dendritic Cell Dichotomy: Identification of IFN- α Producing Cells as a Phenotypically and Functionally Distinct Subset. [J Immunol. 186: 1477-85.](#)
4. Dellacasa-Lindberg, I. *et al.* (2011) Migratory Activation of Primary Cortical Microglia upon Infection with *Toxoplasma gondii*. [Infect Immun. 79: 3046-52.](#)
5. Krejsek, J. *et al.* (2010) Expression of CD200/CD200R regulatory molecules on granulocytes and monocytes is modulated by cardiac surgical operation. [Perfusion 25: 389-97.](#)
6. Rijkers, E.S. *et al.* (2008) The inhibitory CD200R is differentially expressed on human and mouse T and B lymphocytes. [Mol Immunol. 45: 1126-35.](#)
7. Costello, D.A. *et al.* (2011) Long term potentiation is impaired in membrane glycoprotein CD200-deficient mice: a role for Toll-like receptor activation. [J Biol Chem. 286 \(40\): 34722-32.](#)
8. Simelyte, E. *et al.* (2008) CD200-Fc, a novel antiarthritic biologic agent that targets proinflammatory cytokine expression in the joints of mice with collagen-induced arthritis. [Arthritis Rheum. 58: 1038-43.](#)
9. Whitmore, L.C. *et al.* (2014) NOX2 protects against progressive lung injury and multiple organ dysfunction syndrome. [Am J Physiol Lung Cell Mol Physiol. 307: L71-82.](#)
10. Chitnis, T. *et al.* (2007) Elevated neuronal expression of CD200 protects Wlds mice from inflammation-mediated neurodegeneration. [Am J Pathol. 170: 1695-712.](#)
11. Larson, D. *et al.* (2012) Chronic helminth infection reduces basophil responsiveness in an IL-10-dependent manner. [J Immunol. 188: 4188-99.](#)
12. Torrero, M.N. *et al.* (2009) CD200R surface expression as a marker of murine basophil activation. [Clin Exp Allergy. 39 \(3\): 361-9.](#)
13. Hernangómez M *et al.* (2012) CD200-CD200R1 interaction contributes to neuroprotective effects of anandamide on experimentally induced inflammation. [Glia 60 \(9\): 1437-50.](#)
14. Bain, C.C. & Mowat, A.M. (2012) CD200 receptor and macrophage function in the intestine. [Immunobiology 217 \(6\): 643-51.](#)
15. Seeds, R.E. *et al.* (2011) The role of myeloid receptors on murine plasmacytoid dendritic cells in induction of type I interferon. [Int Immunopharmacol. 11 \(7\): 794-801.](#)

16. Akkaya, M. *et al.* (2013) Dissection of agonistic and blocking effects of CD200 receptor antibodies. [PLoS One 8 \(5\): e63325.](#)
17. Cassard L *et al.* (2012) Fcγ receptors inhibit mouse and human basophil activation. [J Immunol. 189 \(6\): 2995-3006.](#)
18. Liu, J.Q. *et al.* (2016) A Critical Role for CD200R Signaling in Limiting the Growth and Metastasis of CD200+ Melanoma. [J Immunol. Jul 6. pii: 1600052. \[Epub ahead of print\]](#)
19. Toya, E.E. & Ohba, M. (2015) Improved Long-term Culture of Epidermal Stem Cells Utilizing CD200R-expressing Feeder Cells [Showa Uni J Med Sci 27 \(2\) 83-91](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®549 , DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	--	------------------	--	---------------	--

'M315591:180503'

Printed on 05 May 2018