

Datasheet: MCA2235BT

Description:	RAT ANTI MOUSE CD206:Biotin
Specificity:	CD206
Other names:	MANNOSE RECEPTOR C TYPE 1
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	MR5D3
Isotype:	IgG2a
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use their own system using appropriate negative/positive controls.

(1) **CD206 is expressed weakly at the cell surface. Staining may be increased following membrane permeabilisation. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

Target Species	Mouse
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Chimaeric CRD4-7-Fc protein
External Database Links	UniProt: Q61830 Related reagents

Entrez Gene:

[17533](#) Mrc1 [Related reagents](#)

Fusion Partners Spleen cells from immunised Fischer rats were fused with cells of the Y3 myeloma cell line

Specificity **Rat anti mouse CD206 antibody, clone MR5D3** recognizes the mouse mannose receptor, a ~175 kDa type 1 membrane glycoprotein that is also known as CD206. CD206 is expressed on most tissue macrophages, certain endothelial cells and *in vitro* derived dendritic cells ([Zamze et al. 2002](#)).

The mannose receptor, CD206, is composed of a N-terminal cysteine-rich domain, a fibronectin type II domain, eight tandemly arranged C-type lectin domains (CTLD), a transmembrane domain, and a cytoplasmic domain. The terminal cysteine-rich domain binds sulfated sugars, and the CTLD recognizes carbohydrates terminating in mannose, fucose and N-acetylglucosamine, all sugars found on microorganisms and on some endogenous proteins ([Su et al. 2005](#)).

Rat anti mouse CD206 antibody, clone MR5D3 has been reported to be non-inhibitory for the binding of the mannose receptor to carbohydrate ligands ([Zamze et al. 2002](#)). Clone MR5D3 has also been shown to work in Western Blot ([Martinez-Pomares et al. 2003](#) and [Su et al. 2005](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Martinez-Pomares, L. *et al.* (2003) Analysis of mannose receptor regulation by IL-4, IL-10, and proteolytic processing using novel monoclonal antibodies. [J Leukoc Biol. 73 \(5\): 604-13.](#)
2. Zamze, S. *et al.* (2002) Recognition of bacterial capsular polysaccharides and lipopolysaccharides by the macrophage mannose receptor. [J Biol Chem. 277 \(44\): 41613-23.](#)
3. Hassan, M.F. *et al.* (2006) The *Schistosoma mansoni* hepatic egg granuloma provides a favorable microenvironment for sustained growth of *Leishmania donovani*. [Am J Pathol. 169: 943-53.](#)
4. Hardison, S.E. *et al.* (2010) Interleukin-17 Is Not Required for Classical Macrophage Activation in a Pulmonary Mouse Model of *Cryptococcus neoformans* Infection. [Infect Immun. 78: 5341-51.](#)
5. Geier, H. & Celli, J. (2011) Phagocytic receptors dictate phagosomal escape and intracellular proliferation of *Francisella tularensis*. [Infect Immun. 79 \(6\): 2204-14.](#)
6. Bacci, M. *et al.* (2009) Macrophages are alternatively activated in patients with endometriosis and required for growth and vascularization of lesions in a mouse model of disease. [Am J Pathol. 175: 547-56.](#)
7. Chavele, K.M. *et al.* (2010) Mannose receptor interacts with Fc receptors and is critical for the development of crescentic glomerulonephritis in mice. [J Clin Invest. 120: 1469-78.](#)
8. deSchoolmeester, M.L. *et al.* (2009) The mannose receptor binds *Trichuris muris* excretory/secretory proteins but is not essential for protective immunity. [Immunology 126: 246-55.](#)
9. Devey, L. *et al.* (2009) Tissue-resident macrophages protect the liver from ischemia reperfusion injury via a heme oxygenase-1-dependent mechanism. [Mol Ther. 17: 65-72.](#)
10. Dewals, B.G. *et al.* (2010) IL-4/Ralpha-independent expression of mannose receptor and Ym1 by macrophages depends on their IL-10 responsiveness. [PLoS Negl Trop Dis. 4 \(5\): e689.](#)
11. Hardison, S.E. *et al.* (2010) Pulmonary infection with an interferon-gamma-producing *Cryptococcus neoformans* strain results in classical macrophage activation and protection. [Am J Pathol. 176: 774-85.](#)
12. Hawkes, C.A. *et al.* (2009) Selective targeting of perivascular macrophages for clearance of beta-amyloid in cerebral amyloid angiopathy. [Proc Natl Acad Sci USA 106: 1261-6.](#)

13. Zehner, M. *et al.* (2011) Mannose receptor polyubiquitination regulates endosomal recruitment of p97 and cytosolic antigen translocation for cross-presentation. [Proc Natl Acad Sci USA 108: 9933-8.](#)
14. Famulski, K.S. *et al.* (2010) Alternative macrophage activation-associated transcripts in T-cell-mediated rejection of mouse kidney allografts. [Am J Transplant 10 \(3\): 490-7.](#)
15. Takagi, H. *et al.* (2009) Cooperation of specific ICAM-3 grabbing nonintegrin-related 1 (SIGNR1) and complement receptor type 3 (CR3) in the uptake of oligomannose-coated liposomes by macrophages. [Glycobiology 19: 258-66.](#)
16. Deepe, G.S. Jr. & Buesing, W.R. (2011) Deciphering the Pathways of Death of *Histoplasma capsulatum*-Infected Macrophages: Implications for the Immunopathogenesis of Early Infection. [J Immunol. 188: 334-44.](#)
17. Schneider, D. *et al.* (2012) Neonatal rhinovirus infection induces mucous metaplasia and airways hyperresponsiveness. [J Immunol. 188 \(6\): 2894-904.](#)
18. Kondo, Y. *et al.* (2011) Macrophages counteract demyelination in a mouse model of globoid cell leukodystrophy. [J Neurosci. 31: 3610-24.](#)
19. Joyce, K.L. *et al.* (2012) Using eggs from *Schistosoma mansoni* as an *in vivo* model of helminth-induced lung inflammation. [J Vis Exp. Jun 5 \(64\): e3905.](#)
20. Su, Y. *et al.* (2005) Glycosylation influences the lectin activities of the macrophage mannose receptor. [J Biol Chem. 280: 32811-20.](#)
21. Verheijden, S. *et al.* (2015) Identification of a chronic non-neurodegenerative microglia activation state in a mouse model of peroxisomal β -oxidation deficiency. [Glia. 63 \(9\): 1606-20.](#)
22. O'Flaherty, B.M. *et al.* (2015) CD8+ T Cell Response to Gammaherpesvirus Infection Mediates Inflammation and Fibrosis in Interferon Gamma Receptor-Deficient Mice. [PLoS One. 10 \(8\): e0135719.](#)
23. Eßlinger M *et al.* (2016) Schizophrenia associated sensory gating deficits develop after adolescent microglia activation. [Brain Behav Immun. May 25. pii: S0889-1591\(16\): 30139-8. \[Epub ahead of print\]](#)
24. Øie, C.I. *et al.* (2016) FITC Conjugation Markedly Enhances Hepatic Clearance of N-Formyl Peptides. [PLoS One. 11 \(8\): e0160602.](#)
25. Manning, C.N. *et al.* (2015) Adipose-derived mesenchymal stromal cells modulate tendon fibroblast responses to macrophage-induced inflammation *in vitro*. [Stem Cell Res Ther. 6: 74.](#)
26. Malada-Edelstein, Y.F. *et al.* (2017) Regulatory role of cytosolic phospholipase A₂ alpha in the induction of CD40 in microglia. [J Neuroinflammation. 14 \(1\): 33.](#)
27. Braune, J. *et al.* (2017) IL-6 Regulates M2 Polarization and Local Proliferation of Adipose Tissue Macrophages in Obesity. [J Immunol. Feb 13. pii: 1600476. \[Epub ahead of print\]](#)
28. Bongiorno, E.K. *et al.* (2017) Type 1 Immune Mechanisms Driven by the Response to Infection with Attenuated Rabies Virus Result in Changes in the Immune Bias of the Tumor Microenvironment and Necrosis of Mouse GL261 Brain Tumors. [J Immunol. May 1. pii: 1601444. \[Epub ahead of print\]](#)
29. Litvack ML *et al.* (2016) Alveolar-like Stem Cell-derived Myb(-) Macrophages Promote Recovery and Survival in Airway Disease. [Am J Respir Crit Care Med. 193 \(11\): 1219-29.](#)
30. Sameshima, A. *et al.* (2015) Teneligliptin improves metabolic abnormalities in a mouse model of postmenopausal obesity. [J Endocrinol. 227 \(1\): 25-36.](#)
31. Eskilsson, A. *et al.* (2014) Distribution of microsomal prostaglandin E synthase-1 in the mouse brain. [J Comp Neurol. 522 \(14\): 3229-44.](#)
32. Hosono, K. *et al.* (2016) Signaling of Prostaglandin E Receptors, EP3 and EP4 Facilitates Wound Healing and Lymphangiogenesis with Enhanced Recruitment of M2 Macrophages in Mice. [PLoS One. 11 \(10\): e0162532.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M246075:140324'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)