

Datasheet: MCA2245PE

Description:	RAT ANTI MOUSE CD41:RPE
Specificity:	CD41
Other names:	INTEGRIN ALPHA IIB
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	MWReg30
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1%	Bovine Serum Albumin	
	5%	Sucrose	
Immunogen	Purified murine platelets		
External Database Links	UniProt: Q9QUM0 Related reagents		

Entrez Gene:

[16399](#) Itga2b [Related reagents](#)

Specificity

Rat anti Mouse CD41 antibody, clone MWReg30 recognizes the mouse integrin alpha IIb subunit CD41. CD41 is a ~125 kDa single pass type 1 transmembrane glycoprotein expressed by platelets, megakaryocytes ([Zhang et al. 2007](#)), mast cells ([Berlanga et al. 2005](#)) and hematopoietic progenitors ([Mitjavila-Garcia et al. 2002](#)). CD41 forms a heterodimer with [CD61](#). The CD41/CD61 complex is important for platelet adhesion and aggregation ([Patel et al. 2003](#)) acting as a receptor for many extracellular matrix proteins including fibronectin, thrombospondin and vitronectin ([Weisel et al. 1992](#)).

Rat anti mouse CD41, clone MWReg30 has been reported to inhibit PMA induced aggregation *in vitro* and to induce hypothermia *in vivo* ([Nieswandt et al. 1999](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Winter, O. *et al.* (2010) Megakaryocytes constitute a functional component of a plasma cell niche in the bone marrow. [Blood. 116: 1867-75.](#)
2. Tamagawa-Mineoka, R. *et al.* (2007) The role of platelets in leukocyte recruitment in chronic contact hypersensitivity induced by repeated elicitation. [Am J Pathol. 170: 2019-29.](#)
3. Takayama, M. *et al.* (2010) Genetic analysis of hierarchical regulation for Gata1 and NF-E2 p45 gene expression in megakaryopoiesis. [Mol Cell Biol. 30: 2668-80.](#)
4. Larson, M.K. and Watson, S.P. (2006) Regulation of proplatelet formation and platelet release by integrin alpha IIb beta3. [Blood. 108: 1509-14.](#)
5. Zanzinger, K. *et al.* (2009) Regulation of triggering receptor expressed on myeloid cells 1 expression on mouse inflammatory monocytes. [Immunology. 128: 185-95.](#)
6. Lutskiy, M.I. *et al.* (2007) WASP localizes to the membrane skeleton of platelets. [Br J Haematol. 139: 98-105.](#)
7. Sullivan, B.P. *et al.* (2010) Protective and damaging effects of platelets in acute cholestatic liver injury revealed by depletion and inhibition strategies. [Toxicol Sci. 115: 286-94.](#)
8. Fujita, R. *et al.* (2013) NF-E2 p45 Is Important for Establishing Normal Function of Platelets. [Mol Cell Biol. 33: 2659-70.](#)
9. Perez, L.E. *et al.* (2008) SH2-inositol phosphatase 1 negatively influences early megakaryocyte progenitors. [PLoS One. 3: e3565.](#)
10. Teeling, J.L. *et al.* (2012) Intracerebral immune complex formation induces inflammation in the brain that depends on Fc receptor interaction [Acta Neuropathol. 124: 479-90.](#)
11. Motohashi, H. *et al.* (2010) NF-E2 domination over Nrf2 promotes ROS accumulation and megakaryocytic maturation. [Blood. 115 \(3\): 677-86.](#)
12. Flierl, U. *et al.* (2015) Phosphorothioate backbone modifications of nucleotide-based drugs are potent platelet activators. [J Exp Med. 212 \(2\): 129-37.](#)
13. Devanathan, V. *et al.* (2015) Platelet Gi protein Gai2 is an essential mediator of thrombo-inflammatory organ damage in mice. [Proc Natl Acad Sci U S A. 112 \(20\): 6491-6.](#)
14. Kamat, V. *et al.* (2015) Microfluidic assessment of functional culture-derived platelets in human thrombi under flow. [Exp Hematol. 43 \(10\): 891-900.e4.](#)
15. Goggs, R. *et al.* (2013) The small GTPase Rif is dispensable for platelet filopodia generation in mice. [PLoS One. 8 \(1\): e54663.](#)
16. Williams CM *et al.* (2015) Identification of roles for the SNARE-associated protein, SNAP29, in mouse platelets. [Nov 20. \[Epub ahead of print\]](#)
17. Cuccurullo, A. *et al.* (2016) Blockade of Thrombopoietin Reduces Organ Damage in

Experimental Endotoxemia and Polymicrobial Sepsis. [PLoS One. 11 \(3\): e0151088.](#)

18. Criel, M. *et al.* (2016) Absence of Pear1 does not affect murine platelet function *in vivo*. [Thromb Res. 146: 76-83.](#)

19. Ryan, J. *et al.* (2016) Myeloid cell-mediated renal injury in rapidly progressive glomerulonephritis depends upon spleen tyrosine kinase. [J Pathol. 238 \(1\): 10-20.](#)

20. Woods SJ *et al.* (2015) Kinetic profiling of *in vivo* lung cellular inflammatory responses to mechanical ventilation. [Am J Physiol Lung Cell Mol Physiol. 308 \(9\): L912-21.](#)

21. Asai, J. *et al.* (2016) Platelets Regulate the Migration of Keratinocytes via Podoplanin/CLEC-2 Signaling during Cutaneous Wound Healing in Mice. [Am J Pathol. 186 \(1\): 101-8.](#)

22. Thomson, A.K. *et al.* (2017) Survival of motor neurone protein is required for normal postnatal development of the spleen. [J Anat. 230 \(2\): 337-46.](#)

Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M300872:170106'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)