

Datasheet: MCA2299FB

Description:	HAMSTER ANTI MOUSE CD61:FITC
Specificity:	CD61
Other names:	INTEGRIN BETA 3 CHAIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	HM beta 3.1
Isotype:	IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Species Cross Reactivity	Reacts with: Rat N.B. Antibody reactivity and working conditions may vary between species.						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation (nm)</th> <th>Emission (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>520</td> </tr> </tbody> </table>	Fluorophore	Excitation (nm)	Emission (nm)	FITC	490	520
Fluorophore	Excitation (nm)	Emission (nm)					
FITC	490	520					
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml						
Immunogen	Mouse alpha 5 beta 3 protein purified from the mouse hybridoma 2B4.						

**External Database
Links**

UniProt:

[O54890](#) [Related reagents](#)

Entrez Gene:

[16416](#) Itgb3 [Related reagents](#)

Fusion Partners

Spleen cells from immunised Armenian hamsters were fused with cells of the P3U1 mouse myeloma cell line.

Specificity

Hamster anti Mouse CD61 antibody, clone HM beta 3-1 recognizes the murine integrin beta 3 subunit (CD61), a ~90 kDa a type I membrane protein, expressed primarily on megakaryocytes, platelets, monocytes, macrophages, granulocytes and endothelial cells. CD61 associates with either the alpha IIb integrin (CD41) or the alpha V integrin (CD51) to form the platelet glycoprotein complex IIb/IIIa and the vitronectin receptor (VNR) respectively. The heterodimers formed with CD61 are receptor for a variety of ligands including fibrinogen, fibronectin, von Willebrands factor (vWF), vitronectin and thrombospondin.

Hamster anti Mouse CD61 antibody, clone HM beta 3-1 is reported to partially inhibit the binding of CD61 to fibronectin ([Yasuda et al. 1995](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.

The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR ([BUF041A/B](#)).

References

1. Yasuda, M. *et al.* (1995) Expression and function of fibronectin binding integrins on rat mast cells. [Int. Immunol. 7:251-258.](#)
2. Hodkinson, P.S. *et al.* (2007) Mammalian NOTCH-1 activates beta1 integrins via the small GTPase R-Ras. [J Biol Chem. 282: 28991-9001.](#)
3. Moore, S.F. *et al.* (2015) Loss of the insulin receptor in murine megakaryocytes/platelets causes thrombocytosis and alterations in IGF signalling. [Cardiovasc Res. 107 \(1\): 9-19.](#)
4. Kraft, S. *et al.* (2016) Identification and characterization of a unique role for EDB fibronectin in phagocytosis. [J Mol Med \(Berl\). 94 \(5\): 567-81.](#)
5. Raouf, J. *et al.* (2016) mPGES-1 deletion affects platelet functions in mice. [Clin Sci \(Lond\). Oct 7. pii: CS20160463. \[Epub ahead of print\]](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody.

Shelf Life

18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet Documentation #10041 available at:
<https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL:FITC \(MCA2356F\)](#)

**North & South
America** Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M306783:170628'

Printed on 28 Jun 2017

© 2017 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)