

## Datasheet: MCA2216PE

<b>Description:</b>	MOUSE ANTI SHEEP CD8:RPE
<b>Specificity:</b>	CD8
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	38.65
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	100 TESTS

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Sheep								
<b>Species Cross Reactivity</b>	Reacts with: Bovine, Goat <b>N.B.</b> Antibody reactivity and working conditions may vary between species.								
<b>Product Form</b>	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized								
<b>Reconstitution</b>	Reconstitute with 1 ml distilled water								
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>RPE 488nm laser</td> <td>496</td> <td>578</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	RPE 488nm laser	496	578		
Fluorophore	Excitation Max (nm)	Emission Max (nm)							
RPE 488nm laser	496	578							
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant								
<b>Buffer Solution</b>	Phosphate buffered saline								
<b>Preservative Stabilisers</b>	0.09% Sodium Azide 1% Bovine Serum Albumin 5% Sucrose								
<b>Immunogen</b>	Ovine efferent lymphocytes.								
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS-1 myeloma cell line.								

<b>Specificity</b>	<p><b>Mouse anti Sheep CD8 antibody, clone 38.65</b> recognizes the ovine CD8 cell surface antigen, which is expressed by the cytotoxic/suppressor subset of T lymphocytes.</p> <p>Under reducing conditions, the antigens immunoprecipitated by Mouse anti Sheep CD8 antibody, clone 38.65 migrate at ~33 kDa and ~36 kDa.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Maddox, J.F. <i>et al.</i> (1985) Surface antigens, SBU-T4 and SBU-T8, of sheep T lymphocyte subsets defined by monoclonal antibodies. <a href="#">Immunology. 55 (4): 739-48.</a></li> <li>Mackay, C.R. <i>et al.</i> (1986) Three distinct subpopulations of sheep T lymphocytes. <a href="#">Eur J Immunol. 16 (1): 19-25.</a></li> <li>Mackay, C.R. <i>et al.</i> (1987) A monoclonal antibody to the p220 component of sheep LCA identifies B cells and a unique lymphocyte subset. <a href="#">Cell Immunol. 110 (1): 46-55.</a></li> <li>Mackay, C.R. <i>et al.</i> (1989) Gamma/delta T cells express a unique surface molecule appearing late during thymic development. <a href="#">Eur J Immunol. 19 (8): 1477-83.</a></li> <li>Mackay, C.R. <i>et al.</i> (1986) Thymocyte subpopulations during early fetal development in sheep. <a href="#">J Immunol. 136 (5): 1592-9.</a></li> <li>Breugelmans, S. <i>et al.</i> (2010) Immunoassay of lymphocyte subsets in ovine palatine tonsils. <a href="#">Acta Histochem. 113(4):416-22</a></li> <li>Lybeck, K.R. <i>et al.</i> (2009) Neutralization of interleukin-10 from CD14(+) monocytes enhances gamma interferon production in peripheral blood mononuclear cells from Mycobacterium avium subsp. paratuberculosis-infected goats. <a href="#">Clin Vaccine Immunol. 16 (7): 1003-11.</a></li> <li>Chan, S.S. <i>et al.</i> (2002) Generation and characterization of ovine dendritic cells derived from peripheral blood monocytes. <a href="#">Immunology. 107: 366-72.</a></li> <li>Davies, M.L. <i>et al.</i> (2004) Architecture of secondary lymphoid tissue in sheep experimentally challenged with scrapie. <a href="#">Immunology. 111: 230-6.</a></li> <li>Elh mouzi-Younes, J. <i>et al.</i> (2010) Ovine CD16+/CD14- blood lymphocytes present all the major characteristics of natural killer cells. <a href="#">Vet Res. 41: 4.</a></li> <li>Lybeck, K.R. <i>et al.</i> (2009) Neutralization of interleukin-10 from CD14(+) monocytes enhances gamma interferon production in peripheral blood mononuclear cells from Mycobacterium avium subsp. paratuberculosis-infected goats. <a href="#">Clin Vaccine Immunol. 16: 1003-11.</a></li> <li>Kallapur, S.G. <i>et al.</i> (2011) Pulmonary and Systemic Inflammatory Responses to Intraamniotic IL-1 alpha in fetal sheep. <a href="#">Am J Physiol Lung Cell Mol Physiol. 301(3):L285-95</a></li> <li>Bruce, C.J. <i>et al.</i> (1999) Depletion of bovine CD8+ T cells with chCC63, a chimaeric mouse-bovine antibody. <a href="#">Vet Immunol Immunopathol. 71 (3-4): 215-31.</a></li> <li>Nfon, C.K. <i>et al.</i> (2012) Innate Immune Response to Rift Valley Fever Virus in Goats. <a href="#">PLoS Negl Trop Dis. 6 (4): e1623.</a></li> <li>Lybeck, K.R. <i>et al.</i> (2012) Intestinal Strictures, Fibrous Adhesions and High Local Interleukin-10 Levels in Goats Infected Naturally with <i>Mycobacterium avium</i> subsp. paratuberculosis. <a href="#">J Comp Pathol. 148: 157-72.</a></li> <li>Olsen, L. <i>et al.</i> (2015) The early intestinal immune response in experimental neonatal ovine cryptosporidiosis is characterized by an increased frequency of perforin expressing NCR1(+) NK cells and by NCR1(-) CD8(+) cell recruitment. <a href="#">Vet Res. 46: 28.</a></li> <li>Goh, S. <i>et al.</i> (2016) Identification of <i>Theileria lestoquardi</i> Antigens Recognized by CD8+ T Cells. <a href="#">PLoS One. 11 (9): e0162571.</a></li> <li>Arranz-Solis, D. <i>et al.</i> (2016) Systemic and local immune responses in sheep after <i>Neospora caninum</i> experimental infection at early, mid and late gestation. <a href="#">Vet Res. 47: 2.</a></li> </ol>
<b>Storage</b>	<p>Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.</p> <p>DO NOT FREEZE.</p>

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Shelf Life</b>	12 months from date of reconstitution.
-------------------	--

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10075 available at: 10075: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:RPE \(MCA929PE\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M301000:170106'

**Printed on 01 Aug 2018**

---

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)