

Datasheet: MCA2356A647

| Description: | HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:Alexa Fluor® 647 |
|---------------|--|
| Specificity: | ARMENIAN HAMSTER IgG NEGATIVE CONTROL |
| Format: | ALEXA FLUOR® 647 |
| Product Type: | Negative/Isotype Control |
| Isotype: | IgG |
| Quantity: | 100 TESTS/1ml |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution | | |
|---|-----|----|----------------|--------------------|--|--|
| Flow Cytometry | • | | | * | | |
| Where this patient, has not been tested for use in a particular technique this does not recognitive | | | | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. *It is recommended that the user titrates the antibody for use in their own system to a concentration equivalent to their test reagent.

| Target Species | Negative Control | | | | | |
|-----------------------------------|--|---------------------|-------------------|--|--|--|
| Product Form | Purified IgG conjug | | | | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) | | | |
| | Alexa Fluor®647 | 650 | 665 | | | |
| Preparation | Purified IgG prepared by ion exchange chromatography | | | | | |
| Buffer Solution | Phosphate buffered saline | | | | | |
| Preservative | 0.09% Sodium Azid | de | | | | |
| Stabilisers | 1% Bovine Seru | ım Albumin | | | | |
| Approx. Protein Concentrations | IgG concentration 0.05 mg/ml | | | | | |
| Immunogen | Trinitrophenol - KLH. | | | | | |
| Specificity | Hamster (Armenian) IgG negative control) is suitable for use as a negative control for the measurement of non-specific binding of Armenian hamster monoclonal antibodies, of isotype to human, mouse or rat tissues. | | | | | |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. | | | | | |
| References | 1. Suzuki-Inoue, K. et al. (2010) Essential in vivo roles of the C-type lectin receptor CLEC-2: | | | | | |

Embryonic/neonatal lethality of CLEC-2 deficient mice by blood/lymp[hatic misconnections and impaired thrombus formation of CLEC-2 deficient platelets. J Biol Chem. 285: 24494-507.

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Europe

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:

10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M300162:170105'

Printed on 01 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint