

Datasheet: MCA2365EL

Description:	MOUSE ANTI BOVINE CD335:Low Endotoxin
Specificity:	CD335
Other names:	NKp46
Format:	Low Endotoxin
Product Type:	Monoclonal Antibody
Clone:	AKS1
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/5 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			
Functional Assays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system to a concentration equivalent to their test reagent.

Target Species	Bovine
Species Cross Reactivity	Reacts with: American Bison N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Endotoxin Level	<0.01EU/ug

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Fusion protein consisting of the extracellular region of bovine CD335.
External Database Links	<p>UniProt: Q863H2 Related reagents</p> <p>Entrez Gene: 369024 NCR1 Related reagents</p>
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the NS-0 myeloma cell line.
Specificity	<p>Mouse anti Bovine CD335 antibody, clone AKS1 recognizes bovine CD335, also known as NKp46 and Natural cytotoxicity triggering receptor 1. CD335 is a type I transmembrane protein, with two extracellular C2-type immunoglobulin-like domains, which functions as an activating receptor. CD335 is expressed by human natural killer cells (Sivori et al. 1997). The bovine homologue is expressed on bovine NK cells (Storset et al. 2004) and no expression of CD335 has been detected on B cells, T cells, monocytes or granulocytes.</p> <p>Clone AKS1 is reported to activate lysis of FcγR-expressing cell line P815, by IL-2 activated NKp46+ cells (Storset et al. 2004).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Storset, A.K. <i>et al.</i> (2004) NKp46 defines a subset of bovine leukocytes with natural killer cell characteristics. Eur J Immunol. 34 (3): 669-76. 2. Kulberg, S. <i>et al.</i> (2004) Reference values for relative numbers of natural killer cells in cattle blood. Dev Comp Immunol. 28 (9): 941-8. 3. Van Rhijn, I. <i>et al.</i> (2007) Massive, sustained γδ T cell migration from the bovine skin <i>in vivo</i>. J Leukoc Biol. 81: 968-73. 4. Toka, F.N. <i>et al.</i> (2011) Rapid and Transient Activation of γδ T Cells to IFN-γ Production, NK Cell-Like Killing, and Antigen Processing during Acute Virus Infection. J Immunol. 186: 4853-61. 5. Bastos, R.G. <i>et al.</i> (2008) Bovine NK cells acquire cytotoxic activity and produce IFN-γ after stimulation by <i>Mycobacterium bovis</i> BCG- or <i>Babesia bovis</i>-exposed splenic dendritic cells. Vet Immunol Immunopathol. 124: 302-12. 6. Boysen P <i>et al.</i> (2006) Bovine CD2-/NKp46+ cells are fully functional natural killer cells with a high activation status. BMC Immunol. 7: 10. 7. Elh mouzi-Younes, J. <i>et al.</i> (2009) Bovine neonate natural killer cells are fully functional and highly responsive to interleukin-15 and to NKp46 receptor stimulation. Vet Res. 40: 54. 8. Elh mouzi-Younes, J. <i>et al.</i> (2010) Ovine CD16+/CD14- blood lymphocytes present all the major characteristics of natural killer cells. Vet Res. 41:4. 9. Hoek, A. <i>et al.</i> (2009) Subpopulations of bovine WC1(+) γδ T cells rather than CD4(+)CD25(high) Foxp3(+) T cells act as immune regulatory cells <i>ex vivo</i>. Vet Res. 40:6. 10. Zuerner, R.L. <i>et al.</i> (2011) A <i>Leptospira borgpetersenii</i> Serovar Hardjo Vaccine Induces a Th1 Response, Activates NK Cells, and Reduces Renal Colonization. Clin Vaccine Immunol. 18: 684-91. 11. Nelson, D.D. <i>et al.</i> (2010) CD8(+)/perforin(+)/WC1(-) γδT cells, not CD8(+) α/β T cells, infiltrate vasculitis lesions of American bison (<i>Bison bison</i>) with experimental sheep-associated malignant catarrhal fever. Vet Immunol Immunopathol. 136: 284-91. 12. Banos, G. <i>et al.</i> (2013) Identification of immune traits correlated with dairy cow health, reproduction and productivity. PLoS One. 8: e65766.

13. Ramstead, A.G. *et al.* (2015) Aging influences the response of T cells to stimulation by the ellagitannin, oenothein B. [Int Immunopharmacol. 26 \(2\): 367-77.](#)
14. Boysen, P. *et al.* (2008) Natural killer cells in lymph nodes of healthy calves express CD16 and show both cytotoxic and cytokine-producing properties. [Dev Comp Immunol. 32 \(7\): 773-83.](#)
15. González-Cano, P. *et al.* (2014) Two functionally distinct myeloid dendritic cell subpopulations are present in bovine blood. [Dev Comp Immunol. 44 \(2\): 378-88.](#)
16. Di Palma, S. *et al.* (2012) Comparative spatiotemporal analysis of the intrathecal immune response in natural listeric rhombencephalitis of cattle and small ruminants. [Comp Immunol Microbiol Infect Dis. 35 \(5\): 429-41.](#)
17. Pirson, C. *et al.* (2015) Highly purified mycobacterial phosphatidylinositol mannosides drive cell-mediated responses and activate NKT cells in cattle. [Clin Vaccine Immunol. 22 \(2\): 178-84.](#)
18. Schneider, D.A. *et al.* (2011) Dynamics of bovine spleen cell populations during the acute response to *Babesia bovis* infection: an immunohistological study. [Parasite Immunol. 33 \(1\): 34-44.](#)
19. Van Meulder F *et al.* (2015) Analysis of the protective immune response following intramuscular vaccination of calves against the intestinal parasite *Cooperia oncophora*. [Int J Parasitol. 45 \(9-10\): 637-46.](#)
20. Patch, J.R. *et al.* (2014) Infection with foot-and-mouth disease virus (FMDV) induces a natural killer (NK) cell response in cattle that is lacking following vaccination. [Comp Immunol Microbiol Infect Dis. 37 \(4\): 249-57.](#)
21. Hedges, J.F. *et al.* (2015) Amphotericin B stimulates $\gamma\delta$ T and NK cells, and enhances protection from *Salmonella* infection. [Innate Immun. 21 \(6\): 598-608.](#)
22. Allan, A.J. *et al.* (2015) Cattle NK Cell Heterogeneity and the Influence of MHC Class I. [J Immunol. 195 \(5\): 2199-206.](#)
23. Maślanka, T. *et al.* (2016) Prostaglandin E₂ exerts the proapoptotic and antiproliferative effects on bovine NK cells [Res Vet Sci. May 25 \[Epub ahead of print\]](#)
24. Hecker YP *et al.* (2015) Cell mediated immune responses in the placenta following challenge of vaccinated pregnant heifers with *Neospora caninum*. [Vet Parasitol. 214 \(3-4\): 247-54.](#)
25. Ohira K *et al.* (2016) Bovine leukemia virus reduces anti-viral cytokine activities and NK cytotoxicity by inducing TGF- β secretion from regulatory T cells. [Immun Inflamm Dis. 4 \(1\): 52-63.](#)
26. Johnson, W.C. *et al.* (2008) Bovine WC1(-) $\gamma\delta$ T cells incubated with IL-15 express the natural cytotoxicity receptor CD335 (NKp46) and produce IFN- γ in response to exogenous IL-12 and IL-18. [Dev Comp Immunol. 32 \(8\): 1002-10.](#)
27. Sipka, A. *et al.* (2016) Bovine natural killer cells are present in *Escherichia coli* infected mammary gland tissue and show antimicrobial activity *in vitro*. [Comp Immunol, Microbiol Infect Dis. Aug 7 \[Epub ahead of print\]](#)
28. Hamilton, C.A. *et al.* (2016) Interactions between natural killer cells and dendritic cells favour T helper1-type responses to BCG in calves. [Vet Res. 47 \(1\): 85.](#)
29. Krueger, L.A. *et al.* (2016) Gamma delta T cells are early responders to *Mycobacterium avium* ssp. *paratuberculosis* in colostrum-replete Holstein calves. [J Dairy Sci. Sep 7. pii: S0022-0302\(16\)30611-7. \[Epub ahead of print\]](#)
30. Steinbach, S. *et al.* (2016) CD4+ and $\gamma\delta$ T Cells are the main Producers of IL-22 and IL-17A in Lymphocytes from *Mycobacterium bovis*-infected Cattle. [Sci Rep. 6: 29990.](#)
31. Rutigliano, H.M. *et al.* (2016) Trophoblast Major Histocompatibility Complex Class I Expression Is Associated with Immune-Mediated Rejection of Bovine Fetuses Produced by Cloning. [Biol Reprod. 95 \(2\): 39.](#)
32. González-Hernández A *et al.* (2016) Host protective ASP-based vaccine against the parasitic nematode *Ostertagia ostertagi* triggers NK cell activation and mixed IgG1-IgG2 response. [Sci Rep. 6: 29496.](#)

Further Reading

1. Sivori, S. *et al.* (1997) p46, a novel natural killer cell-specific surface molecule that mediates cell activation. [J Exp Med. 186 \(7\): 1129-36.](#)
2. Storset, A.K. *et al.* (2003) Natural killer cell receptors in cattle: a bovine killer cell immunoglobulin-like receptor multigene family contains members with divergent signaling motifs.

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10162 available at:
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

- Goat Anti Mouse IgG (STAR76...) [RPE](#)
- Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
- Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@549](#),
[DyLight@649](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)
- Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
- Goat Anti Mouse IgG (STAR77...) [HRP](#)
- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
- Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
- Goat Anti Mouse IgG (STAR70...) [FITC](#)
- Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
- Human Anti Mouse IgG1 (HCA036...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Low Endotoxin \(MCA928EL\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M315655:180503'

Printed on 05 May 2018