

## Datasheet: MCA2385F

<b>Description:</b>	MOUSE ANTI HORSE CD8:FITC
<b>Specificity:</b>	CD8
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	CVS8
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunofluorescence			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Horse								
<b>Species Cross Reactivity</b>	Reacts with: Ass Does not react with: Zebra <b>N.B.</b> Antibody reactivity and working conditions may vary between species.								
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid								
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525		
Fluorophore	Excitation Max (nm)	Emission Max (nm)							
FITC	490	525							
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant								
<b>Buffer Solution</b>	Phosphate buffered saline								
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin								
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1 mg/ml								
<b>Immunogen</b>	Equine PBMC								

**Fusion Partners** Spleen cells from immunised BALB/c mice were fused with cells of the X63-Ag8.653 myeloma cell line.

**Specificity** **Mouse anti Horse CD8 (clone CVS8)**, is a monoclonal antibody recognising the equine homologue of the human CD8 cell surface antigen which is expressed by a subset of T lymphocytes.

A study undertaken using CVS8 to identify CD8 on several wild african equid species indicates that the CVS8 clone recognizes Somali wild ass (*Equus asinus*) but not Grévy's Zebra (*E. grevyi*) or Hartmann's Mountain Zebra (*E. zebra*) ([Ibrahim 2007](#)).

**Flow Cytometry** Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

- References**
1. Lunn, D.P et al (1991) Three monoclonal antibodies identifying antigens on all equine T lymphocytes, and two mutually exclusive T-lymphocyte subsets [Immunology 74: 251-257](#).
  2. Jacks, S. (2007) Experimental infection of neonatal foals with *Rhodococcus equi* triggers adult-like gamma interferon induction. [Clin Vaccine Immunol. 14: 669-77](#).
  3. Pearson, W. *et al.* (2007) Low-dose ginseng (*Panax quinquefolium*) modulates the course and magnitude of the antibody response to vaccination against equid herpesvirus I in horses. [Can J Vet Res. 71: 213-7](#).
  4. Lunn, D.P. *et al.* (1998) Report of the second equine leucocyte antigen workshop, Squaw Valley, California July 1995. [Vet Immunol Immunopathol. 62: 101-143](#).
  5. Merant, C. *et al.* (2003) Cross-species reactivity of seven monoclonal antibodies with equine lymphocytes by flow cytometry. [Vet Res. 34: 791-801](#).
  6. Ibrahim, S (2007) Analysis of monoclonal antibody cross-reactivity with leukocytes from equids and cloning of CD28 [Chapter 5 in PhD Thesis Freie Universität Berlin](#)

**Storage** Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. This product is photosensitive and should be protected from light.  
Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

**Shelf Life** 18 months from date of despatch.

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at:  
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

**Regulatory** For research purposes only

## Related Products

### Recommended Useful Reagents

[MOUSE ANTI HORSE CD4:RPE \(MCA1078PE\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M301767:170109'

Printed on 05 May 2018

