

Datasheet: MCA2352A647T

Description:	HAMSTER ANTI MOUSE CD121a:Alexa Fluor® 647
Specificity:	CD121a
Other names:	INTERLEUKIN 1 RECEPTOR TYPE 1
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	JAMA-147
Isotype:	IgG
Quantity:	25 TESTS/0.25ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10
Functional Assays (1)			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) **Bio-Rad recommend the use of [MCA2352XZ](#) for functional assays.**

Target Species	Mouse						
Product Form	Purified IgG conjugated to Alexa Fluor® 647- liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>Alexa Fluor®647</td> <td>650</td> <td>665</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	Alexa Fluor®647	650	665
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
Alexa Fluor®647	650	665					
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml						
Immunogen	Extracellular domain of murine IL-1R.						
External Database Links	UniProt: P13504 Related reagents						

Entrez Gene:

[16177](#) Il1r1 [Related reagents](#)

Synonyms	Il-1r1, Il1ra
-----------------	---------------

Fusion Partners	Spleen cells from immunised Armenian Hamster were fused with cells of the mouse P3X63.Ag 8-653 myeloma cell line.
------------------------	---

Specificity	<p>Hamster anti Mouse CD121a monoclonal antibody, clone JAMA-147 recognizes murine CD121a, the ~80 kDa type I interleukin-1 receptor. CD121a is a single pass type I membrane glycoprotein, weakly expressed on most cells and tissues including T lymphocytes, thymocytes, epithelial cells and dendritic cells, with stronger expression reported on fibroblasts.</p> <p>The ligands for CD121a are the interleukin-1 (IL-1) family members; IL-1 alpha; IL-1 beta and IL-1 receptor antagonist. IL-1 is a mediator of inflammation and can elicit a variety of other biological responses through binding to CD121a.</p> <p>Hamster anti mouse CD121 antibody, clone JAMA-147 has been reported to block IL-1 dependent cellular responses (Rogers et al. 1992).</p>
--------------------	--

Flow Cytometry	<p>Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
-----------------------	--

References	<ol style="list-style-type: none">1. Wolyniec, K. <i>et al.</i> (2012) E6AP ubiquitin ligase regulates PML-induced senescence in Myc-driven lymphomagenesis Blood. 120: 822-322. Scarim, A.L. <i>et al.</i> (1997) Evidence for the presence of type I IL-1 receptors on beta-cells of islets of Langerhans. Biochim Biophys Acta. 1361 (3): 313-20.3. Merrick, J.C. <i>et al.</i> (1997) Lymphocyte apoptosis during early phase of Listeria infection in mice. Am J Pathol. 151 (3): 785-92.4. Beckerman, K.P. <i>et al.</i> (1993) Release of nitric oxide during the T cell-independent pathway of macrophage activation. Its role in resistance to <i>Listeria monocytogenes</i>. J Immunol. 150 (3): 888-95.5. Kavanagh, D.P. <i>et al.</i> (2013) Mechanisms of adhesion and subsequent actions of a haematopoietic stem cell line, HPC-7, in the injured murine intestinal microcirculation <i>in vivo</i>. PLoS One. 8 (3): e59150.
-------------------	--

Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Acknowledgements	This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic
-------------------------	--

purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA2356A647\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M300200:170105'

Printed on 16 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)