

Datasheet: MCA2395T

Description:	RAT ANTI MOUSE CD13
Specificity:	CD13
Other names:	AMINOPEPTIDASE N
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ER-BMDM1
Isotype:	IgG2a
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Balb/c bone marrow derived macrophages.

**External Database
Links**

UniProt:

[P97449](#) [Related reagents](#)

Entrez Gene:

[16790](#) Anpep [Related reagents](#)

Synonyms

Lap1, Lap-1

Fusion Partners

Cells from immunised rats were fused with cells of the rat Y3-Ag1.2.3 myeloma cell line.

Specificity

Rat anti Mouse CD13 antibody, clone ER-BMDM1 recognizes the murine homologue of human CD13, a ~160 kDa cell surface antigen with aminopeptidase N activity.

In steady state mice, the CD13 antigen is expressed primarily by dendritic cell and mature macrophage subpopulations. The level of CD13 antigen expression increases with maturation, after the monocytic stage. Bone marrow cells and peripheral blood monocytes are reported to be CD13 negative while thioglycollate-elicited macrophages express high levels of CD13.

Dendritic cells, interdigitating dendritic cells, veiled cells and Langerhans cells all express CD13, which is reported to be involved in the development of immature dendritic cells from monocytes.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Leenen, P.J. *et al.* (1992) The monoclonal antibody ER-BMDM1 recognizes a macrophage and dendritic cell differentiation antigen with aminopeptidase activity. [Eur J Immunol. 22 \(6\): 1567-72.](#)
2. Leenen, P.J. *et al.* (1998) Heterogeneity of mouse spleen dendritic cells: in vivo phagocytic activity, expression of macrophage markers, and subpopulation turnover. [J Immunol. 160 \(5\): 2166-73.](#)
3. Sen, A. *et al.* (2012) Innate immune response to homologous rotavirus infection in the small intestinal villous epithelium at single-cell resolution. [Proc Natl Acad Sci U S A. 109: 20667-72.](#)

Further Reading

1. van der Velden, V.H. *et al.* (2001) CD13/aminopeptidase N involvement in dendritic cell maturation. [Leukemia. 15 \(1\): 190-1.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M319118:180719'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)