

Datasheet: MCA2305A488

Description:	RAT ANTI MOUSE CD16/CD32:Alexa Fluor® 488
Specificity:	CD16/CD32
Other names:	MOUSE SEROBLOCK FcR
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	FCR4G8
Isotype:	IgG2b
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Alexa Fluor® 488 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®488	495	519
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	PU5 1.8 IOE7 Balb/c mouse cell line.		
External Database Links	UniProt: P08508 Related reagents P08101 Related reagents		

Entrez Gene:[14131](#) Fcgr3 [Related reagents](#)[14130](#) Fcgr2b [Related reagents](#)

Synonyms	Fcgr2b, Ly-17
Fusion Partners	Cells from immunised rats were fused with cells of the mouse P3X63-Ag8.653 myeloma cell line.
Specificity	Rat anti Mouse CD16/CD32 antibody, clone FCR4G8 recognizes an epitope expressed by the murine low affinity Fc receptors, Fc gamma III (CD16) and Fc gamma II (CD32). In the mouse only the transmembrane form of CD16 is reported to exist, which is expressed on macrophages, NK cells, granulocytes, myeloid precursors, and subpopulations of T lymphocytes. CD32 is primarily expressed on cells of the myeloid lineage and also on mature B lymphocytes.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. da Silva, R.G. <i>et al.</i> (2010) Endothelial alpha3beta1-integrin represses pathological angiogenesis and sustains endothelial-VEGF. Am J Pathol. 177 (3): 1534-48. 2. Birjandi, S.Z. <i>et al.</i> (2011) Alterations in marginal zone macrophages and marginal zone B cells in old mice. J Immunol. 186: 3441-51. 3. Jones, D.T. <i>et al.</i> (2013) Endogenous ribosomal protein L29 (RPL29): a newly identified regulator of angiogenesis in mice. Dis Model Mech. 6: 115-24. 4. Kapellos, T.S. <i>et al.</i> (2016) A Novel Real Time Imaging Platform to Quantify Macrophage Phagocytosis. Biochem Pharmacol. Jul 27. pii: S0006-2952(16)30176-9. [Epub ahead of print] 5. Lunnon, K. <i>et al.</i> (2011) Systemic inflammation modulates Fc receptor expression on microglia during chronic neurodegeneration. J Immunol. 186 (12): 7215-24. 6. Teeling, J.L. <i>et al.</i> (2012) Intracerebral immune complex formation induces inflammation in the brain that depends on Fc receptor interaction. Acta Neuropathol. 124 (4): 479-90. 7. Murinello, S. <i>et al.</i> (2014) Fcγ receptor upregulation is associated with immune complex inflammation in the mouse retina and early age-related macular degeneration. Invest Ophthalmol Vis Sci. 55 (1): 247-58. 8. Hart, A.D. <i>et al.</i> (2012) Age related changes in microglial phenotype vary between CNS regions: grey versus white matter differences. Brain Behav Immun. 26 (5): 754-65. 9. Birjandi, S.Z. <i>et al.</i> (2011) Alterations in marginal zone macrophages and marginal zone B cells in old mice. J Immunol. 186 (6): 3441-51. 10. Hart, R. & Greaves, D.R. (2010) Chemerin contributes to inflammation by promoting macrophage adhesion to VCAM-1 and fibronectin through clustering of VLA-4 and VLA-5. J Immunol. 185 (6): 3728-39.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Acknowledgements	This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in

research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M299884:170105'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)