

Datasheet: MCA2386

Description:	RAT ANTI MOUSE CD223
Specificity:	CD223
Other names:	LAG-3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	C9B7W
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/50 - 1/200
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Murine CD223 Ig fusion protein.

**External Database
Links**

UniProt:

[Q61790](#) [Related reagents](#)

Entrez Gene:

[16768](#) Lag3 [Related reagents](#)

Fusion Partners

Cells from immunised Lewis rats were fused with cells of the Sp/20 myeloma cell line.

Specificity

Rat anti Mouse CD223 antibody, clone C9B7W recognizes murine lymphocyte activation gene-3 (LAG-3), a ~70 kDa activation-induced cell surface molecule that is also referred to as CD223.

Murine CD223 is expressed on activated CD4 positive and CD8 positive alpha/beta T lymphocytes and a subset of natural killer (NK) cells. CD223 binds to MHC class II molecules with high affinity and is reported to negatively regulate T cell homeostasis and T cell expansion.

Clone C9B7W recognizes an epitope within the D2 domain of CD223. Rat anti Mouse CD223 antibody, clone C9B7W is reported to block the *in vitro* function of murine LAG-3 but does not block binding of LAG-3 to MHC class II ([Workman *et al.* 2002](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Workman, C.J. *et al.* (2002) Phenotypic analysis of the murine CD4-related glycoprotein, CD223 (LAG-3). [Eur J Immunol. 32 \(8\): 2255-63.](#)
2. Workman, C.J. & Vignali, D.A. (2005) Negative regulation of T cell homeostasis by lymphocyte activation gene-3 (CD223). [J Immunol. 174 \(2\): 688-95.](#)
3. Byrne, K.T. *et al.* (2011) Autoimmune melanocyte destruction is required for robust CD8+ memory T cell responses to mouse melanoma. [J Clin Invest. 121 \(5\): 1797-809.](#)
4. Workman, C.J. *et al.* (2002) Cutting edge: molecular analysis of the negative regulatory function of lymphocyte activation gene-3. [J Immunol. 169 \(10\): 5392-5.](#)
5. Hu, Z. *et al.* (2013) Regulatory CD8+ T cells associated with erosion of immune surveillance in persistent virus infection suppress *in vitro* and have a reversible proliferative defect. [J Immunol. 191 \(1\): 312-22.](#)
6. Iwasaki, Y. *et al.* (2013) Egr-2 transcription factor is required for Blimp-1-mediated IL-10 production in IL-27-stimulated CD4+ T cells. [Eur J Immunol. 43: 1063-73.](#)
7. Ordway, D. *et al.* (2007) The hypervirulent *Mycobacterium tuberculosis* strain HN878 induces a potent TH1 response followed by rapid down-regulation. [J Immunol. 179: 522-31.](#)
8. Woo, S.R. *et al.* (2010) Differential subcellular localization of the regulatory T-cell protein LAG-3 and the coreceptor CD4. [Eur J Immunol. 40: 1768-77.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®549 , DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG1 NEGATIVE CONTROL \(MCA1211\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M315663:180503'

Printed on 05 May 2018