

Datasheet: MCA2393A488T

Description:	RAT ANTI MOUSE ER-HR3:Alexa Fluor® 488
Specificity:	ER-HR3
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	ER-HR3
Isotype:	IgG2c
Quantity:	0.25 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Product Form	Purified IgG conjugated to Alexa Fluor® 488 - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>Alexa Fluor®488</td> <td>495</td> <td>519</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	Alexa Fluor®488	495	519
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
Alexa Fluor®488	495	519					
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative	0.09% Sodium Azide						
Stabilisers	1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml						
Immunogen	Adherent F1 (CBAxBL) bone marrow stromal cells.						
Fusion Partners	Cells from immunised rats were fused with cells of the mouse P3 - X63 - Ag8.563 myeloma cell line.						
Specificity	Rat anti Mouse ER-HR3, clone ER-HR3 recognizes the murine antigen ER-HR3. ER-HR3 is a cell surface antigen expressed by Langerhans cells in epithelium, a subset of mature macrophages and						

dendritic cells located predominantly in haematopoietic and lymphoid organs. ER-HR3 demonstrates very low levels of expression on peripheral blood monocytes.

During foetal development, ER-HR3 positive cells are localised to haemopoietic islands and are often associated with erythroid progenitor cells. The functions of the ER-HR3 antigen have not been established but reports suggest that the antigen may be involved in adult erythropoiesis and in the regulation of the immune response.

Rat anti Mouse ER-HR3, clone ER-HR3 does not inhibit T cell proliferation in antigen-specific T-cell proliferation studies.

Rat anti Mouse ER-HR3, clone ER-HR3 recognizes two proteins of ~69 kDa and a minor one of ~55 kDa under non-reducing conditions ([de Jong et al. 1994](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.

References

1. de Jong, J.P. *et al.* (1994) A monoclonal antibody (ER-HR3) against murine macrophages. I. Ontogeny, distribution and enzyme histochemical characterization of ER-HR3-positive cells. [Cell Tissue Res. 275: 567-76.](#)
2. Oliveira, M.A. *et al.* (2003) Immature macrophages derived from mouse bone marrow produce large amounts of IL-12p40 after LPS stimulation. [J Leukoc Biol. 74: 857-67.](#)
3. Throsby, M. *et al.* (2000) CD11c+ eosinophils in the murine thymus: developmental regulation and recruitment upon MHC class I-restricted thymocyte deletion. [J Immunol. 165:1965-75.](#)
4. Grabbe, S. *et al.* (2002) Beta2 integrins are required for skin homing of primed T cells but not for priming naive T cells. [J Clin Invest. 109: 183-92.](#)
5. Sonoda, Y. and Sasaki, K. (2012) Hepatic extramedullary hematopoiesis and macrophages in the adult mouse: histometrical and immunohistochemical studies. [Cells Tissues Organs. 196: 555-64.](#)
6. Jacobsen, R.N. *et al.* (2014) Mobilization with granulocyte colony-stimulating factor blocks medullar erythropoiesis by depleting F4/80+VCAM1+CD169+ER-HR3+Ly6G+ erythroid island macrophages in the mouse. [Exp Hematol. pii: S0301-472X\(14\)00139-8.](#)
7. Vogel, J. *et al.* (2003) Transgenic mice overexpressing erythropoietin adapt to excessive erythrocytosis by regulating blood viscosity. [Blood. 102 \(6\): 2278-84.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or

not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M312430:180215'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)