

Datasheet: MCA2387A647T

Description:	RAT ANTI MOUSE Gr-1:Alexa Fluor® 647
Specificity:	Gr-1
Other names:	Ly-6G
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	RB6-8C5
Isotype:	IgG2b
Quantity:	25 TESTS/0.25ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Alexa Fluor® 647 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®647	650	665
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		
Immunogen	Normal murine bone marrow cells.		
External Database Links	UniProt: P35461 Related reagents		

Entrez Gene:

[546644](#) Ly6g [Related reagents](#)

Specificity

Rat anti Mouse Gr-1 antibody, clone RB6-8C5 recognizes the mouse Gr-1 antigen, a ~21–25 kDa GPI anchored cell surface protein bearing a single uPAR/Ly6 domain that belongs to the Ly-6 family of proteins ([Lee et al. 2013](#)). Rat anti Mouse Gr-1 antibody, clone RB6-8C5 reacts predominantly with the Ly-6G protein but weaker reactivity with the Ly-6C protein has been reported ([Fleming et al. 1993](#)). However, other observations dispute the cross-reactivity of clone RB6-8C5 with the Ly-6C protein with the alternative explanation that certain sub-populations of bone marrow cells simultaneously express both Ly-6C and Ly-6G ([Nagendra et al. 2007](#))

The Gr-1 antigen is primarily a marker of myeloid differentiation. In the bone marrow the level of Gr-1 expression is low on immature myeloblasts and increases as the myeloid cells mature to granulocytes. Gr-1 is also expressed on macrophages and transiently on differentiating monocytes.

Rat anti Mouse Gr-1 antibody, clone RB6-8C5 has been used successfully for the depletion of mature neutrophils *in vivo* ([Czuprynski, C.J. et al 1994](#), [Daley et al. 2008](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Fleming, T.J. et al. (1993) Selective expression of Ly-6G on myeloid lineage cells in mouse bone marrow. RB6-8C5 mAb to granulocyte-differentiation antigen (Gr-1) detects members of the Ly-6 family. [J Immunol. 151 \(5\): 2399-408.](#)
2. Hestdal, K. et al. (1991) Characterization and regulation of RB6-8C5 antigen expression on murine bone marrow cells. [J Immunol. 147 \(1\): 22-8.](#)
3. Czuprynski, C.J. et al. (1994) Administration of anti-granulocyte mAb RB6-8C5 impairs the resistance of mice to *Listeria monocytogenes* infection. [J Immunol. 152 \(4\): 1836-46.](#)
4. Sumagin R et al. (2010) LFA-1 and Mac-1 define characteristically different intraluminal crawling and emigration patterns for monocytes and neutrophils *in situ*. [J Immunol. 185 \(11\): 7057-66.](#)
5. Takano, K. et al. (2011) Successful treatment of acute lung injury with pitavastatin in septic mice: potential role of glucocorticoid receptor expression in alveolar macrophages. [J Pharmacol Exp Ther. 336: 381-90.](#)
6. Giroux, M. et al. (2011) SMAD3 prevents graft-versus-host disease by restraining Th1 differentiation and granulocyte-mediated tissue damage. [Blood. 117: 1734-44.](#)
7. Suttman, H. et al. (2006) Neutrophil granulocytes are required for effective Bacillus Calmette-Guérin immunotherapy of bladder cancer and orchestrate local immune responses. [Cancer Res. 66: 8250-7.](#)
8. Nix, R.N. et al. (2007) Hemophagocytic macrophages harbor *Salmonella enterica* during persistent infection. [PLoS Pathog. 3: e193.](#)
9. Kanda, N. et al. (2011) Visfatin Enhances CXCL8, CXCL10, and CCL20 Production in Human Keratinocytes. [Endocrinology. 152: 3155-64.](#)
10. Conlan, J. and North, R. (1994) Neutrophils are essential for early anti-*Listeria* defense in the liver, but not in the spleen or peritoneal cavity, as revealed by a granulocyte-depleting monoclonal antibody. [J Exp Med. 179:259-68.](#)
11. Takebe, M. et al. (2014) Inhibition of histone deacetylases protects septic mice from lung and splenic apoptosis. [J Surg Res. 187 \(2\): 559-70.](#)
12. Francke, A. et al. (2011) Generation of mature murine monocytes from heterogeneous bone marrow and description of their properties. [J Histochem Cytochem. 59: 813-25.](#)
13. Sharp, P.E. et al. (2013) FcγRIIb on myeloid cells and intrinsic renal cells rather than B cells protects from nephrotoxic nephritis. [J Immunol. 190: 340-8.](#)
14. Hamers, A.A. et al. (2014) Limited role of nuclear receptor Nur77 in *Escherichia coli*-induced peritonitis. [Infect Immun. 82 \(1\): 253-64.](#)

15. Roche, J.A. *et al.* (2015) Myofiber damage precedes macrophage infiltration after *in vivo* injury in dysferlin-deficient a/j mouse skeletal muscle. [Am J Pathol. 185 \(6\): 1686-98.](#)
16. Lee, Y.S. *et al.* (2015) Interleukin-1 (IL-1) signaling in intestinal stromal cells controls KC/CXCL1 secretion, which correlates with recruitment of IL-22- secreting neutrophils at early stages of *Citrobacter rodentium* infection. [Infect Immun. 83 \(8\): 3257-67.](#)
17. Heckelsmiller, K. *et al.* (2002) Combined dendritic cell- and CpG oligonucleotide-based immune therapy cures large murine tumors that resist chemotherapy. [Eur J Immunol. 32 \(11\): 3235-45.](#)
18. Zhang, M.Z. *et al.* (2015) Inhibition of cyclooxygenase-2 in hematopoietic cells results in salt-sensitive hypertension. [J Clin Invest. 125 \(11\): 4281-94.](#)
19. Leblond, A.L. *et al.* (2015) Systemic and Cardiac Depletion of M2 Macrophage through CSF-1R Signaling Inhibition Alters Cardiac Function Post Myocardial Infarction. [PLoS One. 10 \(9\): e0137515.](#)
20. Kojo, K. *et al.* (2016) BLT1 signalling protects the liver against acetaminophen hepatotoxicity by preventing excessive accumulation of hepatic neutrophils. [Sci Rep. 6: 29650.](#)
21. Otsuru, S. *et al.* (2017) Hematopoietic derived cells do not contribute to osteogenesis as osteoblasts. [Bone. 94: 1-9.](#)
22. Wang, Y. *et al.* (2015) Proximal tubule-derived colony stimulating factor-1 mediates polarization of renal macrophages and dendritic cells, and recovery in acute kidney injury. [Kidney Int. 88 \(6\): 1274-1282.](#)
23. Cousins, F.L. *et al.* (2016) Evidence for a dynamic role for mononuclear phagocytes during endometrial repair and remodelling. [Sci Rep. 6: 36748.](#)
24. Cotrina ML *et al.* (2017) Direct comparison of microglial dynamics and inflammatory profile in photothrombotic and arterial occlusion evoked stroke. [Neuroscience. 343: 483-494.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M300527:170105'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)