

Datasheet: MCA2309

Description:	MOUSE ANTI PIG CD11R3	
Specificity: CD11R3		
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	2F4/11	
Isotype:	IgG1	
Quantity:	0.25 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			1/25 - 1/50
Immunohistology - Frozen	-			
Immunohistology - Paraffin				
ELISA			•	
Immunoprecipitation	-			
Western Blotting	-			
Functional Assays (1)			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Removal of sodium azide is recommended prior to use in functional assays - Bio-Rad recommend the use of $\underline{EQU003}$ for this purpose.

Target Species	Pig
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Porcine alveolar macrophages

Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the X63-Ag.8.653 myeloma cell line
Specificity	Mouse anti Pig CD11R3, clone 2F4/11 recognizes porcine CD11R3, a ~155 kDa cell surface glycoprotein, member of the alpha integrin family. Mouse anti Pig CD11R3, clone 2F4/11 was clustered as CD11R3 at the Third International Workshop on Swine Leukocyte Differentiation Antigens (Haverson et al. 2001). CD11R3 has a similar expression pattern to the human CD11b marker, being expressed on granulocytes, monocytes and alveolar macrophages, but not on lymphocytes, eythrocytes or platelets (Dominguez et al. 2001). Mouse anti Pig CD11R3, clone 2F4/11 is reported to block phagocytosis of complement-opsonized
	zymosan particles by polymorphonuclear granulocytes and alveolar macrophages (<u>Bullido et al.</u> 1996).
Flow Cytometry	Use 10ul of the suggested working dilution to 1x10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Porcine spleen.
Western Blotting	MCA2309 detects a band of approximately 155 kDa in alveolar macrophage lysates under reducing conditions.
References	 Sbrana, S. <i>et al.</i> (2014) Phenotype Changes of Circulating Monocytes in a Hypercholesterolemic Swine Model of Coronary Artery Disease J Cytol Histol 5:270. Domínguez, J. <i>et al.</i> (2001) Workshop studies on monoclonal antibodies in the myeloid panel with CD11 specificity. Vet Immunol Immunopathol. 80 (1-2): 111-9. Sánchez-Torres C <i>et al.</i> (2003) Expression of porcine CD163 on monocytes/macrophages correlates with permissiveness to African swine fever infection. Arch Virol. 148 (12): 2307-23. Van de Walle, G.R. <i>et al.</i> (2003) Transmission of pseudorabies virus from immune-masked blood monocytes to endothelial cells. J Gen Virol. 84 (Pt 3): 629-37. Alvarez, B. <i>et al.</i> (2000) Molecular and functional characterization of porcine LFA-1 using monoclonal antibodies to CD11a and CD18. Xenotransplantation 7: 258-266 Sánchez, C. <i>et al.</i> (1999) The porcine 2A10 antigen is homologous to human CD163 and related to macrophage differentiation. J Immunol. 162: 5230-7 Thorgersen, E.B. <i>et al.</i> (2010) Anti-inflammatory effects of C1-Inhibitor in porcine and human whole blood are independent of its protease inhibition activity. Innate Immun. 16: 254-64 Thorgersen, E.B. <i>et al.</i> (2010) CD14 inhibition efficiently attenuates early inflammatory and hemostatic responses in <i>Escherichia coli</i> sepsis in pigs. FASEB J. 24: 712-22 Baert K <i>et al.</i> (2015) Cell type-specific differences in β-glucan recognition and signalling in porcine innate immune cells. Dev Comp Immunol. 48 (1): 192-203. Barratt-Due, A. <i>et al.</i> (2011) <i>Ornithodoros moubata</i> Complement Inhibitor Is an Equally Effective C5 Inhibitor in Pigs and Humans. J Immunol. 187: 4913-9 Jacobsen, M.J. (2015) Altered Methylation Profile of Lymphocytes Is Concordant with Perturbation of Lipids Metabolism and Inflammatory Response in Obesity Journal of Diabetes Research Article ID 610546 (Epub ahead of print] Crisci, E. <i>et al.</i>

Obese Porcine Model of Metabolic Syndrome. <u>J Diabetes Res. 2016: 3486727.</u>

Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res 39: 54				
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denaturate antibody. Should this product contain a precipitate we recommend microcentrifugation before use.	re			
Shelf Life	18 months from date of despatch.				
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf				
Regulatory	For research purposes only				

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Rabbit Anti Mouse IgG (STAR9...)

Goat Anti Mouse IgG (STAR77...)

Rabbit Anti Mouse IgG (STAR12...)

RPE

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP
Rabbit Anti Mouse IgG (STAR8...) DyLight®800

Goat Anti Mouse IgG (STAR70...) FITC

Rabbit Anti Mouse IgG (STAR13...) HRP

Human Anti Mouse IgG1 (HCA036...) HRP

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®549,

<u>DyLight®649</u>, <u>DyLight®680</u>, <u>DyLight®800</u>,

FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

'M317052:180618'

Printed on 21 Jun 2018

@ 2018 Bio-Rad Laboratories Inc $\mid \underline{\text{Legal}} \mid \underline{\text{Imprint}}$