

Datasheet: MCA2484

Description:	MOUSE ANTI HUMAN CD282
Specificity:	CD282
Other names:	TLR2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	11G7
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/100 - 1/200
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation	■			
Western Blotting			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	UniProt:

Entrez Gene:

[7097](#) TLR2 [Related reagents](#)

Synonyms	TIL4
Specificity	Mouse anti Human CD282 antibody, clone 11G7 recognizes the human Toll-like receptor 2 (TLR2), a ~90 kDa type 1 transmembrane protein that is primarily expressed by peripheral blood monocytes. TLR2, also known as CD282, is reported to respond to a diverse range of bacterial cell wall components, mediating the innate immune response in co-operation with MD-2. Mouse anti Human CD282 antibody, clone 11G7 has been reported to block IL-6 cytokine production in response to ara-lipoarabinomannan (araLAM) and tripalmitoyl cysteinyl lipopeptide (Pam3CSK4) (Sandor <i>et al.</i> 2003). Clone 11G7 does not inhibit the production of cytokines induced by zymosan stimulation.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	1. Sandor, F. <i>et al.</i> (2003) Importance of extra- and intracellular domains of TLR1 and TLR2 in NFkappa B signaling. J Cell Biol. 162 (6): 1099-110. 2. Zhou, S. <i>et al.</i> (2010) Discovery of a novel TLR2 signaling inhibitor with anti-viral activity. Antiviral Res. 87 (3): 295-306. 3. Royer, P.J. <i>et al.</i> (2010) The mannose receptor mediates the uptake of diverse native allergens by dendritic cells and determines allergen-induced T cell polarization through modulation of IDO activity. J Immunol. 185 (3): 1522-31. 4. Chen, E.S. <i>et al.</i> (2010) Serum amyloid A regulates granulomatous inflammation in sarcoidosis through Toll-like receptor-2. Am J Respir Crit Care Med. 181 (4): 360-73. 5. Nguyen, D.N. <i>et al.</i> (2016) Oral antibiotics increase blood neutrophil maturation and reduce bacteremia and necrotizing enterocolitis in the immediate postnatal period of preterm pigs. Innate Immun. 22 (1): 51-62. 6. Royer, P.J. <i>et al.</i> (2010) The mannose receptor mediates the uptake of diverse native allergens by dendritic cells and determines allergen-induced T cell polarization through modulation of IDO activity. J Immunol. 185 (3): 1522-31.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M315748:180503'

Printed on 21 Jun 2018