

Datasheet: MCA2465F

Description:	RAT ANTI MOUSE CD273:FITC
Specificity:	CD273
Other names:	PD-L2
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	TY25
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	B7-DC transfected RAW264.7 cells.		
External Database Links	UniProt: Q9WUL5 Related reagents		

Entrez Gene:

[58205](#) Pdcd1lg2 [Related reagents](#)

Synonyms	B7dc, Btdc, Cd273, Pdl2
Fusion Partners	Lymph node cells from immunised SD rats were fused with the cells of the P3U1 myeloma cell line.
Specificity	Rat anti Mouse CD273 antibody, clone TY25 recognises mouse CD273, also known as B7-dendritic cell (B7-DC), or programmed death ligand-2 (PD-L2). CD273 is a 247 amino acid, ~42 kDa member of the B7 family. The expression of CD273 is restricted to a subpopulation of dendritic cells (DCs), and DCs and macrophages that have been stimulated with interferon gamma, GM-CSF or IL-4. CD273 has been identified as a ligand for PD-1(CD279), along with another B7 family member B7-H1 (CD274). Studies suggest that both CD273 and B7-H1 co-stimulate or inhibit T cell proliferation and cytokine production. Rat anti Mouse CD273 antibody, clone TY25 is reported to block the binding of CD273 (B7-DC) to it's receptor (Matsumoto et al. 2004). Removal of sodium azide is recommended prior to use in functional assays, the use of EQU003 is recommended for this purpose.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B/C)
References	1. Matsumoto, K. <i>et al.</i> (2004) B7-DC regulates asthmatic response by an IFN-gamma-dependent mechanism. J Immunol. 172 (4): 2530-41. 2. Kanai, T. <i>et al.</i> (2003) Blockade of B7-H1 suppresses the development of chronic intestinal inflammation. J Immunol. 171 (8): 4156-63. 3. Yamazaki, T. <i>et al.</i> (2005) Blockade of B7-H1 on macrophages suppresses CD4+ T cell proliferation by augmenting IFN-gamma-induced nitric oxide production. J Immunol. 175: 1586-92. 4. Furuhashi K <i>et al.</i> (2012) Mouse lung CD103+ and CD11b high dendritic cells preferentially induce distinct CD4+ T-cell responses. Am J Respir Cell Mol Biol. 46 (2): 165-72. 5. López-Medina M <i>et al.</i> (2015) <i>Salmonella</i> induces PD-L1 expression in B cells. Immunol Lett. pii: S0165-2478(15)30018-3. 6. López-Medina M <i>et al.</i> (2015) <i>Salmonella</i> impairs CD8 T cell response through PD-1: PD-L axis. Immunobiology. pii: S0171-2985(15)30028-0.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M301477:170109'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)