

Datasheet: MCA2462T

Description:	RAT ANTI MOUSE CD80
Specificity:	CD80
Other names:	B7-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	RM80
Isotype:	IgG2a
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/50
Immunohistology - Frozen	■			
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	BCL1 cells expressing CD80.

**External Database
Links**

UniProt:

[Q00609](#)

[Related reagents](#)

Entrez Gene:

[12519](#)

Cd80

[Related reagents](#)

Synonyms

B7

Fusion Partners

Spleen cells from immunised SD rats were fused with cells of the P3U1 myeloma cell line.

Specificity

Rat anti Mouse CD80 antibody, clone RM80 recognizes mouse CD80 (B7-1), a ~60 kDa cell surface glycoprotein which is a member of the CD28/B7 family. In mice, CD80 is expressed on monocytes, peritoneal macrophages and dendritic cells, and expression may be significantly increased upon B lymphocytes by LPS and by IL-4.

CD80 has been identified as a ligand for CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA-4), two structurally similar molecules expressed on T cells. CD28 and CTLA4 are two receptors that have essential but opposing functions in T-cell stimulation. The Interaction of CD80 with CD28 stimulates and sustains T cell responses, whereas the interaction of CD80 with CTLA4 is reported to inhibit T-cell responses.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.

References

1. Nuriya, S. *et al.* (1996) The differential role of CD86 and CD80 co-stimulatory molecules in the induction and the effector phases of contact hypersensitivity. [Int Immunol. 8 \(6\): 917-26.](#)
2. Jin LP *et al.* (2004) Adoptive transfer of paternal antigen-hyporesponsive T cells induces maternal tolerance to the allogeneic fetus in abortion-prone matings. [J Immunol. 173 \(6\): 3612-9.](#)
3. Nakajima, A. *et al.* (1997) Requirement of CD28-CD86 co-stimulation in the interaction between antigen-primed T helper type 2 and B cells. [Int Immunol. 9 \(5\): 637-44.](#)
4. Nozawa, K. *et al.* (2001) Preferential blockade of CD8(+) T cell responses by administration of anti-CD137 ligand monoclonal antibody results in differential effect on development of murine acute and chronic graft-versus-host diseases. [J Immunol. 167 \(9\): 4981-6.](#)
5. Bedoret, D. *et al.* (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. [J Clin Invest. 119 \(12\): 3723-38.](#)
6. Legutko, A. *et al.* (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor- γ Activity in Dendritic Cells. [J Immunol. 187: 4517-29.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®549 , DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M315736:180503'

Printed on 05 May 2018