

Datasheet: MCA2463PE

Description:	RAT ANTI MOUSE CD86:RPE
Specificity:	CD86
Other names:	B7-2
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	PO3
Isotype:	IgG2b
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1.0 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1%	Bovine Serum Albumin	
	5%	Sucrose	
Immunogen	Mouse B-cell line, BCL1.		
External Database Links	UniProt: P42082 Related reagents		

Entrez Gene:[12524](#) Cd86 [Related reagents](#)

Fusion Partners	Spleen cells from immunized Sprague Dawley rats were fused with cells of the P3U1 myeloma cell line.
Specificity	Rat anti mouse CD86 antibody, clone PO3 recognizes mouse CD86 (B7-2), a ~80 kDa cell surface glycoprotein, a member of the CD28/B7 family. In mouse CD86 is expressed at high levels on peripheral blood monocytes and dendritic cells, and at low levels on resting B and T-lymphocytes. Expression of CD86, on these cell populations, can be increased upon activation. CD86 has been identified, along with CD80 (B7-1), as a ligand for CD28 and cytotoxic T-lymphocyte antigen-4 (CTLA4). CD28 and CTLA4 are two receptors that have opposing functions in T-cell stimulation. Interaction of CD86 with CD28 promotes a number of T-cell activities, whereas the binding of CD86 to CTLA4 inhibits T-cell responses. Rat anti mouse CD86 antibody, clone PO3 has been reported to block some co-stimulatory functions of CD86 (Nakajima et al. 1997).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Jin, L.P. <i>et al.</i> (2004) Adoptive transfer of paternal antigen-hyporesponsive T cells induces maternal tolerance to the allogeneic fetus in abortion-prone matings. J Immunol. 173 (6): 3612-9. 2. Nozawa, K. <i>et al.</i> (2001) Preferential blockade of CD8(+) T cell responses by administration of anti-CD137 ligand monoclonal antibody results in differential effect on development of murine acute and chronic graft-versus-host diseases. J Immunol. 167 (9): 4981-6. 3. Nakajima, A. <i>et al.</i> (1997) Requirement of CD28-CD86 co-stimulation in the interaction between antigen-primed T helper type 2 and B cells. Int Immunol. 9 (5): 637-44. 4. Bedoret, D. <i>et al.</i> (2009) Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. J Clin Invest. 119 (12): 3723-38. 5. Legutko, A. <i>et al.</i> (2011) Sirtuin 1 Promotes Th2 Responses and Airway Allergy by Repressing Peroxisome Proliferator-Activated Receptor-γ Activity in Dendritic Cells. J Immunol. 187: 4517-29.
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M301030:170106'

Printed on 01 May 2018

