

Datasheet: MCA2548FT

Description:	MOUSE ANTI HUMAN CD142:FITC
Specificity:	CD142
Other names:	TISSUE FACTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	CLB/TF-5
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1mg/ml
External Database Links	UniProt: P13726 Related reagents Entrez Gene: 2152 F3 Related reagents
Specificity	Mouse anti Human CD142 antibody, clone CLB/TF-5 recognizes human CD142, a 45kD cell surface glycoprotein which is otherwise known as Tissue Factor (TF). CD142 expression can be

induced on monocytes, macrophages and endothelial cells by various stimuli including interleukin 1, tumour necrosis factor and endotoxin. CD142 initiates the blood clotting cascade by binding coagulation factor VIIa, which activates factor IX or factor X by specific limited proteolysis. CD142 also plays an important role in inflammation, angiogenesis, and the pathophysiology of atherosclerosis and cancer.

Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
-----------------------	---

References	1. VanDerPutten, R.F. <i>et al.</i> (2005) High-affinity antibodies in a new immunoassay for plasma tissue factor: reduction in apparent intra-individual variation. Clin Chem Lab Med. 43 (12): 1386-91. 2. Mobarrez, F. <i>et al.</i> (2010) A multicolor flow cytometric assay for measurement of platelet-derived microparticles. Thromb Res. 125: e110-6. 3. Tehrani, S. <i>et al.</i> (2010) Atorvastatin has antithrombotic effects in patients with type 1 diabetes and dyslipidemia. Thromb Res. 126: e225-31. 4. Aass, H.C. <i>et al.</i> (2011) Fluorescent particles in the antibody solution result in false TF- and CD14-positive microparticles in flow cytometric analysis. Cytometry A. 79: 990-9. 5. Trummer, A. <i>et al.</i> (2008) Isotype controls in phenotyping and quantification of microparticles: a major source of error and how to evade it. Thromb Res. 122: 691-700.
-------------------	--

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	--

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation available at: Material Safety Datasheet Documentation #10058 available at: https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M301307:170109'

Printed on 23 Aug 2017