

Datasheet: MCA2537B

Description:	MOUSE ANTI HUMAN CD16:Biotin
Specificity:	CD16
Other names:	FcRIII
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	DJ130c
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Macaque N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG conjugated to Biotin - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)
Stabilisers	1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1mg/ml
External Database Links	<p>UniProt:</p> <p>P08637 Related reagents</p> <p>O75015 Related reagents</p> <p>Entrez Gene:</p>

[2214](#) FCGR3A [Related reagents](#)

[2215](#) FCGR3B [Related reagents](#)

Synonyms CD16A, CD16B, FCG3, FCGR3, IGFR3

Specificity **Mouse anti Human CD16 antibody, clone DJ130c** recognizes human CD16, also known as Low affinity immunoglobulin gamma Fc region receptor III-A or Fc-gamma RIIIa. CD16a is a 254 amino acid ~50-65 kDa single pass type 1 transmembrane glycoprotein bearing two [Ig-like C2 type](#) domains. CD16 exists as a transmembranous form (Fc gammaRIIIA, or CD16A) and a glycosyl phosphatidylinositol (GPI) anchored form, Fc gammaRIIIB, or CD16B ([Scallon et al. 1989](#)). CD16A is expressed by NK cells, some T cells, and macrophages, whereas CD16B is primarily expressed by granulocytes ([Ravetch and Perussia 1989](#)). In addition, CD16B exists as two allelic variants NA1 and NA2 . DJ130c recognizes all polymorphonuclear cells irrespective of their NA phenotype.

Mouse anti Human CD16 antibody, clone DJ130c recognizes an epitope in the first membrane-distal domain of CD16, recognizes both CD16a and CD16b and has been demonstrated to cross-react with CD16 from rhesus macaques, *Macaca mulatta* ([Xu et al. 2012](#))

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

- References**
- Schmidt, R.E. (1993) CD16 cluster workshop report. In Leucocyte Typing V: White cell differentiation antigens, Vol.1. Edited by Schlossman, S.F. *et al.* Oxford University Press. p805 – 806.
 - Kakko, T. *et al.* (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. [Diab Vasc Dis Res. 8: 221-8.](#)
 - Shantsila, E. *et al.* (2012) Fibrinolytic status in acute coronary syndromes: evidence of differences in relation to clinical features and pathophysiological pathways. [Thromb Haemost. 108: 32-40.](#)
 - Shantsila, E. *et al.* (2011) Immunophenotypic characterization of human monocyte subsets: possible implications for cardiovascular disease pathophysiology. [J Thromb Haemost. 9: 1056-66.](#)
 - Tapp, L.D. *et al.* (2012) The CD14++CD16+ monocyte subset and monocyte-platelet interactions in patients with ST-elevation myocardial infarction. [J Thromb Haemost. 10: 1231-41.](#)
 - Ambarus, C.A. *et al.* (2012) Intimal lining layer macrophages but not synovial sublining macrophages display an IL-10 polarized-like phenotype in chronic synovitis. [Arthritis Res Ther. 14: R74.](#)
 - Ambarus, C.A. *et al.* (2012) Systematic validation of specific phenotypic markers for in vitro polarized human macrophages. [J Immunol Methods. 375: 196-206.](#)
 - Ambarus, C.A. *et al.* (2012) Soluble immune complexes shift the TLR-induced cytokine production of distinct polarized human macrophage subsets towards IL-10. [PLoS One. 7: e35994.](#)
 - Shantsila, E. *et al.* (2012) The effects of exercise and diurnal variation on monocyte subsets and monocyte-platelet aggregates. [Eur J Clin Invest. 42: 832-9.](#)
 - Chehadeh, W. *et al.* (2009) Antibody-mediated opsonization of red blood cells in parvovirus B19 infection. [Virology. 390: 56-63.](#)
 - Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. [Circ Heart Fail. 6: 127-35.](#)
 - Jaipersad, A.S. *et al.* (2014) Expression of monocyte subsets and angiogenic markers in relation to carotid plaque neovascularization in patients with pre-existing coronary artery disease and carotid stenosis. [Ann Med. 11: 1-9.](#)
 - Shantsila, E. *et al.* (2015) Free Light Chains in patients with acute coronary syndromes: Relationships to inflammation and renal function. [Int J Cardiol. 185: 322-7.](#)

14. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. [Circ Heart Fail. 6 \(1\): 127-35.](#)
15. Romee R *et al.* (2013) NK cell CD16 surface expression and function is regulated by a disintegrin and metalloprotease-17 (ADAM17). [Blood. 121 \(18\): 3599-608.](#)
16. Sousa, S. *et al.* (2015) Human breast cancer cells educate macrophages toward the M2 activation status. [Breast Cancer Res. 17: 101.](#)

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M290829:160715'

Printed on 25 May 2018