

Datasheet: MCA2537T

Description:	MOUSE ANTI HUMAN CD16
Specificity:	CD16
Other names:	FcRIII
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	DJ130c
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/20 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Species Cross Reactivity	Reacts with: Macaque N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	<p>UniProt:</p> <p>P08637 Related reagents</p> <p>O75015 Related reagents</p> <p>Entrez Gene:</p> <p>2214 FCGR3A Related reagents</p> <p>2215 FCGR3B Related reagents</p>
Synonyms	CD16A, CD16B, FCG3, FCGR3, IGFR3
Specificity	<p>Mouse anti Human CD16 antibody, clone DJ130c recognizes human CD16, also known as Low affinity immunoglobulin gamma Fc region receptor III-A or Fc-gamma RIIIa. CD16a is a 254 amino acid ~50-65 kDa single pass type 1 transmembrane glycoprotein bearing two Ig-like C2 type domains. CD16 exists as a transmembranous form (Fc gammaRIIIA, or CD16A) and a glycosyl phosphatidylinositol (GPI) anchored form, Fc gammaRIIIB, or CD16B (Scallon et al. 1989). CD16A is expressed by NK cells, some T cells, and macrophages, whereas CD16B is primarily expressed by granulocytes (Ravetch and Perussia 1989). In addition, CD16B exists as two allelic variants NA1 and NA2 . DJ130c recognizes all polymorphonuclear cells irrespective of their NA phenotype.</p> <p>Mouse anti Human CD16 antibody, clone DJ130c recognizes an epitope in the first membrane-distal domain of CD16, recognizes both CD16a and CD16b and has been demonstrated to cross-react with CD16 from rhesus macaques, <i>Macaca mulatta</i> (Xu et al. 2012)</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Human tonsil
References	<ol style="list-style-type: none"> Schmidt, R.E. (1993) CD16 cluster workshop report. In Leucocyte Typing V: White cell differentiation antigens, Vol.1. Edited by Schlossman, S.F. <i>et al.</i> Oxford University Press. p805 – 806. Kakko, T. <i>et al.</i> (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. Diab Vasc Dis Res. 8: 221-8. Shantsila, E. <i>et al.</i> (2012) Fibrinolytic status in acute coronary syndromes: evidence of differences in relation to clinical features and pathophysiological pathways. Thromb Haemost. 108: 32-40. Shantsila, E. <i>et al.</i> (2011) Immunophenotypic characterization of human monocyte subsets: possible implications for cardiovascular disease pathophysiology. J Thromb Haemost. 9: 1056-66. Tapp, L.D. <i>et al.</i> (2012) The CD14++CD16+ monocyte subset and monocyte-platelet interactions in patients with ST-elevation myocardial infarction. J Thromb Haemost. 10: 1231-41. Ambarus, C.A. <i>et al.</i> (2012) Intimal lining layer macrophages but not synovial sublining macrophages display an IL-10 polarized-like phenotype in chronic synovitis. Arthritis Res Ther. 14: R74. Ambarus, C.A. <i>et al.</i> (2012) Systematic validation of specific phenotypic markers for in vitro polarized human macrophages. J Immunol Methods. 375: 196-206. Ambarus, C.A. <i>et al.</i> (2012) Soluble immune complexes shift the TLR-induced cytokine

- production of distinct polarized human macrophage subsets towards IL-10. [PLoS One. 7: e35994.](#)
9. Shantsila, E. *et al.* (2012) The effects of exercise and diurnal variation on monocyte subsets and monocyte-platelet aggregates. [Eur J Clin Invest. 42: 832-9.](#)
10. Chehadah, W. *et al.* (2009) Antibody-mediated opsonization of red blood cells in parvovirus B19 infection. [Virology. 390: 56-63.](#)
11. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. [Circ Heart Fail. 6: 127-35.](#)
12. Jaipersad, A.S. *et al.* (2014) Expression of monocyte subsets and angiogenic markers in relation to carotid plaque neovascularization in patients with pre-existing coronary artery disease and carotid stenosis. [Ann Med. 11: 1-9.](#)
13. Shantsila, E. *et al.* (2015) Free Light Chains in patients with acute coronary syndromes: Relationships to inflammation and renal function. [Int J Cardiol. 185: 322-7.](#)
14. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. [Circ Heart Fail. 6 \(1\): 127-35.](#)
15. Romee R *et al.* (2013) NK cell CD16 surface expression and function is regulated by a disintegrin and metalloprotease-17 (ADAM17). [Blood. 121 \(18\): 3599-608.](#)
16. Sousa, S. *et al.* (2015) Human breast cancer cells educate macrophages toward the M2 activation status. [Breast Cancer Res. 17: 101.](#)

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@549 , DyLight@649 , DyLight@680 , DyLight@800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com
'M315783:180503'

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

Printed on 25 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)