

Datasheet: MCA2516PE

Description:	MOUSE ANTI HUMAN CD87:RPE
Specificity:	CD87
Other names:	UPAR, UPA-R
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	VIM-5
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1.0ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
	5% Sucrose		
Immunogen	THP-1 cells (Human acute monocytic leukaemia cells).		
External Database Links	UniProt:		
	Q03405	Related reagents	

Entrez Gene:

[5329](#) PLAUR [Related reagents](#)

Synonyms	MO3, UPAR
-----------------	-----------

Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.
------------------------	---

Specificity	<p>Mouse anti Human CD87 antibody, clone VIM-5 recognizes human CD87, also known as urokinase plasminogen activator receptor (uPAR), a 50-65 kDa single chain glycosylphosphatidylinositol (GPI) anchored glycoprotein, expressed by granulocytes, tissue macrophages, monocytes, endothelial cells, fibroblasts and mesenchymal and epithelial tumour cells.</p> <p>CD87 specifically binds and activates urokinase plasminogen activator (uPA), a serine protease which converts the inactive receptor-bound proenzyme plasminogen into plasmin, triggering the proteolysis cascade of fibrinolysis. The interaction between uPAR and uPA is also known to play a role in cancer cell invasion and metastasis.</p> <p>Clone VIM-5 is reported to block urokinase binding.</p>
--------------------	--

Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
-----------------------	---

References	<ol style="list-style-type: none">1. Sillaber, C. <i>et al.</i> (1997) Molecular and functional characterization of the urokinase receptor on human mast cells. J Biol Chem. 272 (12): 7824-32.2. Gadd, S.J. <i>et al.</i> (1990) M5, a phosphoinositol-linked human myelomonocytic activation-associated antigen. Clin Exp Immunol. 80 (2): 252-6.
-------------------	--

Further Reading	<ol style="list-style-type: none">1. Mondino, A. & Blasi, F. (2004) uPA and uPAR in fibrinolysis, immunity and pathology. Trends Immunol. 25 (8): 450-5.
------------------------	--

Storage	<p>Prior to reconstitution store at +4°C.</p> <p>After reconstitution store at +4°C.</p> <p>DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	---

Shelf Life	12 months from date of reconstitution.
-------------------	--

Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M312225:180209'

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)