

Datasheet: MCA2628F

Description:	MOUSE ANTI HUMAN CD279:FITC
Specificity:	CD279
Other names:	PD-1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MIH4
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
Immunogen	Human CD279 - transfected L cells.		
External Database Links	UniProt: Q15116 Related reagents		

Entrez Gene:[5133](#) PDCD1 [Related reagents](#)

Synonyms	PD1
Fusion Partners	Spleen cells from immunised C3H mice were fused with cells of the P3U1 myeloma cell line.
Specificity	<p>Mouse anti Human CD279 antibody, clone MIH4 detects CD279, a co-stimulatory molecule also known as programmed cell death-1 (PD-1). CD279 is a ~50-55 kDa membrane protein which is a member of the CD28 family, and functions mainly as a negative regulator of T-cell activation. CD279 has two specific ligands; CD274 (PD-L1) and CD273 (PD-L2), and their interaction is key in the balance between stimulatory and inhibitory signals needed for effective immune responses to microbes and self-tolerance. CD279 is inducibly expressed by T-cells, B-cells, NK-T-cells and monocytes upon activation.</p> <p>Loss of CD279 function has been associated with a number of autoimmune diseases, including rheumatoid arthritis, type I diabetes and ankylosing spondylitis. Recent studies suggest that CD279 could be targeted therapeutically in the treatment of HIV infection to reduce T-cell exhaustion.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	1. Kanai, T. <i>et al.</i> (2003) Blockade of B7-H1 suppresses the development of chronic intestinal inflammation. J Immunol. 171 (8): 4156-63.
Further Reading	1. Freeman, G.J. <i>et al.</i> (2006) Reinvigorating exhausted HIV-specific T cells via PD-1-PD-1 ligand blockade. J Exp Med. 203 (10): 2223-7. 2. Keir, M.E. <i>et al.</i> (2007) PD-1 and its ligands in T-cell immunity. Curr Opin Immunol. 19 (3): 309-14.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com 'M314914:180424'	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

Printed on 27 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)