

Datasheet: MCA2607

Description:	MOUSE ANTI HUMAN iC3b (NEOANTIGEN)
Specificity:	iC3b (NEOANTIGEN)
Other names:	INACTIVATED COMPLEMENT COMPONENT 3b
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	013III-1.16
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			
Functional Assays			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Borate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Purified human iC3b
External Database	UniProt:

Links [P01024](#) [Related reagents](#)

Entrez Gene:

[718](#) C3 [Related reagents](#)

Synonyms CPAMD1

Specificity **Mouse anti Human iC3b (Neoantigen) antibody, clone 013III-1.16** recognizes the ~42 kDa human inactive complement component 3b (iC3b) neoantigen present in blood serum. Cleavage of C3b into iC3b by the protease factor I, in the presence of cofactors, changes the structure and resulting binding properties, preventing binding by complement factor B and properdin. This prevents further progression of the complement pathway, providing another mechanism by which the complement pathway may be regulated.

iC3b is thought to play a role in inducing tolerance, binding complement receptor type 3 on antigen presenting cells and stimulating them to produce transforming growth factor beta2 and interleukin-10. It is thought to stimulate B-cells via the CD21 receptor providing a link between the innate and adaptive immune responses. Additionally iC3b acts as an opsonin which is recognised by complement receptor 3 expressed on leukocytes or complement receptor of the immunoglobulin superfamily (CRIg) on Kupffer cells, stimulating phagocytosis of the tagged cells. iC3b levels are often elevated in diseases such as Systemic Lupus Erythematosis and Rheumatoid arthritis. iC3b levels are often elevated in diseases such as Systemic Lupus Erythematosis and Rheumatoid arthritis.

Histology Positive Control Tissue Kidney from post streptococcal glomerulonephritis patients

Further Reading 1. Janssen, B. *et al.* (2007) Structural insights into the central complement component C3. [Molecular Immunology. 44: 3-10.](#)
2. Bergmann-Leitner, E.S. *et al.* (2006) Complement 3d: from molecular adjuvant to target of immune escape mechanisms. [Clin Immunol. 121 \(2\): 177-85.](#)
3. Sohn, J.H. *et al.* (2003) Tolerance is dependent on complement C3 fragment iC3b binding to antigen-presenting cells. [Nat Med. 9 \(2\): 206-12.](#)

Storage Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet Documentation #10077 available at: <https://www.bio-rad-antibodies.com/uploads/MSDS/10077.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight@800](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@549](#),
[DyLight@649](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Human Anti Mouse IgG2b (HCA038...) [FITC](#), [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)

Recommended Useful Reagents

[HISTAR DETECTION SYSTEM \(STAR3000A\)](#)
[HISTAR DETECTION SYSTEM \(STAR3000B\)](#)
[HISTAR DETECTION SYSTEM \(STAR3000C\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M283688:160308'

Printed on 06 Jun 2017

© 2017 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)