

Datasheet: MCA2748

Description:	RAT ANTI MOUSE CD36
Specificity:	CD36
Other names:	GPIV (IIIb)
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MF3
Isotype:	IgG2a
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/25
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	IL-4 treated murine thioglycollate-elicited peritoneal macrophages

**External Database
Links**

UniProt:

[Q08857](#) [Related reagents](#)

Entrez Gene:

[12491](#) Cd36 [Related reagents](#)

Fusion Partners

Spleen cells from immunised DA rats were fused with cells of the Y3Ag 1.2.3 myeloma cell line

Specificity

Rat anti Mouse CD36 antibody, clone MF3 recognizes mouse CD36, also known as platelet glycoprotein 4, glycoprotein IIIb or PAS IV. CD36 is an ~85 kDa multipass transmembrane glycoprotein primarily expressed on platelets, monocytes/macrophages, smooth muscle and endothelial cells. The CD36 molecule is type B scavenger receptor, which binds to multiple ligands including thrombospondin, anionic phospholipids, oxidized low density lipoproteins and long chain fatty acids.

CD36 has diverse functions and is reported to play a role in innate immunity, platelet adhesion/aggregation and long chain fatty acid transport. The CD36 molecule also directly mediates cytoadhesion of erythrocytes infected with *Plasmodium falciparum*, and may be involved in the development of atherosclerotic lesions and the formation of foam cells.

Rat anti Mouse CD36 antibody, clone MF3 has been shown to inhibit IL-4 induced thioglycollate-elicited peritoneal macrophage fusion and significantly block IL-4/GM-CSF-induced bone-marrow derived macrophage fusion.

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Helming, L. *et al.* (2009) The scavenger receptor CD36 plays a role in cytokine-induced macrophage fusion. [J Cell Sci. 122 \(Pt 4\): 453-9.](#)
2. Mwaikambo, B.R. *et al.* (2009) Hypoxia up-regulates CD36 expression and function via hypoxia-inducible factor-1- and phosphatidylinositol 3-kinase-dependent mechanisms. [J Biol Chem. 284: 26695-707.](#)
3. Yang, C.N. *et al.* (2011) Mechanism mediating oligomeric A β clearance by naïve primary microglia. [Neurobiol Dis. 42 \(3\): 221-30.](#)
4. Seeds, R.E. *et al.* (2011) The role of myeloid receptors on murine plasmacytoid dendritic cells in induction of type I interferon. [Int Immunopharmacol. 11: 794-801.](#)
5. Samovski, D. *et al.* (2012) Insulin and AMPK regulate FA translocase/CD36 plasma membrane recruitment in cardiomyocytes via Rab GAP AS160 and Rab8a Rab GTPase. [J Lipid Res. 53 \(4\): 709-17.](#)
6. Zhou, D. *et al.* (2012) CD36 level and trafficking are determinants of lipolysis in adipocytes. [FASEB J. 26 \(11\): 4733-42.](#)
7. Nie, S. *et al.* (2015) Detection of atherosclerotic lesions and intimal macrophages using CD36-targeted nanovesicles. [J Control Release. pii: S0168-3659\(15\)30173-5.](#)
8. Samovski, D. *et al.* (2015) Regulation of AMPK activation by CD36 links fatty acid uptake to β -oxidation. [Diabetes. 64 \(2\): 353-9.](#)
9. Pugholm LH *et al.* (2015) Enhanced humoral responses induced by targeting of antigen to murine dendritic cells. [Scand J Immunol. Sep 8. \[Epub ahead of print\]](#)
10. Miyazaki H *et al.* (2014) Fatty acid binding protein 7 regulates phagocytosis and cytokine production in Kupffer cells during liver injury. [Am J Pathol. 184 \(9\): 2505-15.](#)
11. Sui, Y. *et al.* (2018) Perinatal Bisphenol A Exposure Increases Atherosclerosis in Adult Male PXR-Humanized Mice. [Endocrinology. 159 \(4\): 1595-608.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
-------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@649 , DyLight@800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M317166:180619'

Printed on 01 Aug 2018