

## Datasheet: MCA275EL

<b>Description:</b>	MOUSE ANTI RAT CD11b:Low Endotoxin
<b>Specificity:</b>	CD11b
<b>Other names:</b>	INTEGRIN ALPHA M CHAIN, MAC-1
<b>Format:</b>	Low Endotoxin
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-42
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.5 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/50 - 1/100
Immunohistology - Paraffin (1)			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative /positive controls.

(1)**OX-42 is reported to be suitable for paraffin-embedded sections following PLP fixation (Whiteland *et al.*, 1995).**

<b>Target Species</b>	Rat
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	None present
<b>Carrier Free</b>	Yes
<b>Endotoxin Level</b>	<0.01 EU/ug

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Resident rat peritoneal macrophages.
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the NSO/U mouse myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Rat CD11b, clone OX-42</b>, recognizes rat CD11b, also known as <a href="#">integrin alpha-M</a>, the receptor for the iC3b component of complement. CD11b is a 1151 amino acid single pass type 1 transmembrane glycoprotein possessing a single <a href="#">vWFA</a> domain and multiple <a href="#">FG-GAP</a> repeats. CD11b is expressed on most macrophages, including resident and activated peritoneal macrophages and Kupffer cells and around 35% of alveolar macrophages. The antibody also labels dendritic cells, granulocytes and <a href="#">microglia</a> in the brain (<a href="#">Robinson et al.1986</a>).</p> <p>Mouse anti Rat CD11b, clone OX-42 is reported to inhibit complement mediated rosettes (<a href="#">Robinson et al.1986</a>) as well as inhibit myelin binding and uptake (<a href="#">van der Laan et al.1996</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood
<b>References</b>	<ol style="list-style-type: none"> <li>1. Robinson, A.P. <i>et al.</i> (1986) Macrophage heterogeneity in the rat as delineated by two monoclonal antibodies MRC OX-41 and MRC OX-42, the latter recognizing complement receptor type 3. <a href="#">Immunology. 57 (2): 239-47.</a></li> <li>2. Barclay, A.N. (1981) The localization of population of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. <a href="#">Immunology. 42: 593-600.</a></li> <li>3. Milligan, C.E. <i>et al.</i> (1991) Differential immunochemical markers reveal the normal distribution of brain macrophages and microglia in the developing rat brain. <a href="#">J Comp Neurol. 314 (1): 125-35.</a></li> <li>4. Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. <a href="#">J Histochem Cytochem. 43 (3): 313-20.</a></li> <li>5. Zilka, N. <i>et al.</i> (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. <a href="#">J Neuroimmunol. 209 (1-2): 16-25.</a></li> <li>6. Shokouhi, B.N. <i>et al</i> (2010) Microglial responses around intrinsic CNS neurons are correlated with axonal regeneration. <a href="#">BMC Neurosci. 11: 13.</a></li> <li>7. Calvo, M. <i>et al.</i> (2010) Neuregulin-ErbB signaling promotes microglial proliferation and chemotaxis contributing to microgliosis and pain after peripheral nerve injury. <a href="#">J Neurosci. 30 (15): 5437-50.</a></li> <li>8. Jean, Y.H. <i>et al.</i> (2009) Capnellene, a natural marine compound derived from soft coral, attenuates chronic constriction injury-induced neuropathic pain in rats <a href="#">Br J Pharmacol. 158: 713-25.</a></li> <li>9. Morales-Garcia, J.A. <i>et al.</i> (2011) Phosphodiesterase 7 inhibition preserves dopaminergic neurons in cellular and rodent models of Parkinson disease. <a href="#">PLoS One. 6(2):e17240.</a></li> <li>10. Spencer-Segal, J.L. <i>et al.</i> (2011) Distribution of Phosphorylated TrkB Receptor in the Mouse Hippocampal Formation Depends on Sex and Estrous Cycle Stage. <a href="#">J Neurosci. 31: 6780-90.</a></li> <li>11. Chew, S.S. <i>et al.</i> (2011) Response of retinal Connexin43 to optic nerve injury. <a href="#">Invest Ophthalmol Vis Sci. 52: 3620-9.</a></li> <li>12. Huh, S.H. <i>et al</i> (2011) Ethyl pyruvate rescues nigrostriatal dopaminergic neurons by regulating glial activation in a mouse model of Parkinson's disease. <a href="#">J Immunol. 187: 960-9.</a></li> <li>13. Jeong, H.K. <i>et al.</i> (2010) Inflammatory responses are not sufficient to cause delayed neuronal death in ATP-induced acute brain injury. <a href="#">PLoS One. 5: e13756.</a></li> <li>14. Liew, H.K. <i>et al.</i> (2012) Systemic administration of urocortin after intracerebral hemorrhage reduces neurological deficits and neuroinflammation in rats. <a href="#">J Neuroinflammation. 9: 13.</a></li> </ol>

15. Szmydynger-Chodobska, J. *et al.* (2011) Multiple sites of vasopressin synthesis in the injured brain. [J Cereb Blood Flow Metab. 31: 47-51.](#)
16. Tchoukalova, Y.D. *et al.* (2012) *In vivo* adipogenesis in rats measured by cell kinetics in adipocytes and plastic-adherent stroma-vascular cells in response to high-fat diet and thiazolidinedione. [Diabetes. 61: 137-44.](#)
17. Xu, Q. *et al.* (2011) Spinal phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling cascades in inflammation-induced hyperalgesia. [J Neurosci. 31: 2113-24.](#)
18. Lovett-Barr, M.R. *et al.* (2012) Repetitive intermittent hypoxia induces respiratory and somatic motor recovery after chronic cervical spinal injury. [J Neurosci. 32 \(11\): 3591-600.](#)
19. Lavis, S. *et al.* (2012) Reactive astrocytes overexpress TSPO and are detected by TSPO positron emission tomography imaging. [J Neurosci. 32: 10809-18.](#)
20. Ortega, F.J. *et al.* (2012) Glibenclamide enhances neurogenesis and improves long-term functional recovery after transient focal cerebral ischemia. [J Cereb Blood Flow Metab. 33: 356-64.](#)
21. Zhao, H. *et al.* (2013) Brain 3-Mercaptopyruvate Sulfurtransferase (3MST): Cellular Localization and Downregulation after Acute Stroke. [PLoS One. 8\(6\):e67322.](#)
22. Huang S *et al.* (2015) Expression of Peroxiredoxin 1 After Traumatic Spinal Cord Injury in Rats. [Cell Mol Neurobiol. May 24. \[Epub ahead of print\]](#)
23. Wang, A.Y. *et al.* (2015) Searching for a rat model of chronic tympanic membrane perforation: Healing delayed by mitomycin C/dexamethasone but not paper implantation or iterative myringotomy. [Int J Pediatr Otorhinolaryngol. 79 \(8\): 1240-7.](#)
24. Hernangómez M *et al.* (2016) CD200R1 agonist attenuates glial activation, inflammatory reactions, and hypersensitivity immediately after its intrathecal application in a rat neuropathic pain model. [J Neuroinflammation. 13 \(1\): 43.](#)
25. Signarovitz, A.L. *et al.* (2012) Mucosal immunization with live attenuated *Francisella novicida* U112ΔigB protects against pulmonary *F. tularensis* SCHU S4 in the Fischer 344 rat model. [PLoS One. 7: e47639.](#)
26. Liu, Z. *et al.* (2016) Leukocyte Infiltration Triggers Seizure Recurrence in a Rat Model of Temporal Lobe Epilepsy. [Inflammation. Apr 4. \[Epub ahead of print\].](#)
27. Szmydynger-Chodobska, J. *et al.* (2016) The Involvement of Pial Microvessels in Leukocyte Invasion after Mild Traumatic Brain Injury. [PLoS One. 11 \(12\): e0167677.](#)
28. Ejaz, S. *et al.* (2014) MRI and neuropathological validations of the involvement of air pollutants in cortical selective neuronal loss. [Environ Sci Pollut Res Int. 21 \(5\): 3351-62.](#)
29. Alizadeh, A. *et al.* (2017) Neuregulin-1 positively modulates glial response and improves neurological recovery following traumatic spinal cord injury. [Glia. Apr 29. \[Epub ahead of print\]](#)
30. Popiolek-Barczyk, K. *et al.* (2017) Biphalin, a Dimeric Enkephalin, Alleviates LPS-Induced Activation in Rat Primary Microglial Cultures in Opioid Receptor-Dependent and Receptor-Independent Manners. [Neural Plasticity. 2017: 1-19.](#)
31. Huang RY *et al.* (2017) Rapid and Delayed Effects of Pulsed Radiofrequency on Neuropathic Pain: Electrophysiological, Molecular, and Behavioral Evidence Supporting Long-Term Depression. [Pain Physician. 20 \(2\): E269-E283.](#)
32. Bourke, G. *et al.* (2017) Effects of early nerve repair on experimental brachial plexus injury in neonatal rats. [J Hand Surg \(Eur Vol\). 1753193417732696 \[Epub ahead of print\].](#)
33. Wang, Z.C. *et al.* (2017) Involvement of NF-κB and the CX3CR1 Signaling Network in Mechanical Allodynia Induced by Tetanic Sciatic Stimulation. [Neurosci Bull. Jun 13 \[Epub ahead of print\].](#)

---

**Storage**

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

<b>Shelf Life</b>	18 months from date of despatch.
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10162 available at: 10162: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight@800</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Human Anti Mouse IgG2a (HCA037...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@549</a> , <a href="#">DyLight@649</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:Low Endotoxin \(MCA1210EL\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M315872:180503'

**Printed on 20 Jun 2018**