

Datasheet: MCA275PE

Description:	MOUSE ANTI RAT CD11b:RPE
Specificity:	CD11b
Other names:	INTEGRIN ALPHA M CHAIN, MAC-1
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	OX-42
Isotype:	IgG2a
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by ion exchange chromatography		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
	5% Sucrose		
Immunogen	Resident rat peritoneal macrophages.		
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NSO/U mouse myeloma cell line.		

Specificity

Mouse anti Rat CD11b, clone OX-42, recognizes rat CD11b, also known as [integrin alpha-M](#), the receptor for the iC3b component of complement. CD11b is a 1151 amino acid single pass type 1 transmembrane glycoprotein possessing a single [vWFA](#) domain and multiple [FG-GAP](#) repeats. CD11b is expressed on most macrophages, including resident and activated peritoneal macrophages and Kupffer cells and around 35% of alveolar macrophages. The antibody also labels dendritic cells, granulocytes and [microglia](#) in the brain ([Robinson et al.1986](#)).

Mouse anti Rat CD11b, clone OX-42 is reported to inhibit complement mediated rosettes ([Robinson et al.1986](#)) as well as inhibit myelin binding and uptake ([van der Laan et al.1996](#)).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Robinson, A.P. *et al.* (1986) Macrophage heterogeneity in the rat as delineated by two monoclonal antibodies MRC OX-41 and MRC OX-42, the latter recognizing complement receptor type 3. [Immunology. 57 \(2\): 239-47.](#)
2. Barclay, A.N. (1981) The localization of population of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. [Immunology. 42: 593-600.](#)
3. Milligan, C.E. *et al.* (1991) Differential immunochemical markers reveal the normal distribution of brain macrophages and microglia in the developing rat brain. [J Comp Neurol. 314 \(1\): 125-35.](#)
4. Whiteland, J.L. *et al.* (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. [J Histochem Cytochem. 43 \(3\): 313-20.](#)
5. Zilka, N. *et al.* (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. [J Neuroimmunol. 209 \(1-2\): 16-25.](#)
6. Shokouhi, B.N. *et al* (2010) Microglial responses around intrinsic CNS neurons are correlated with axonal regeneration. [BMC Neurosci. 11: 13.](#)
7. Calvo, M. *et al.* (2010) Neuregulin-ErbB signaling promotes microglial proliferation and chemotaxis contributing to microgliosis and pain after peripheral nerve injury. [J Neurosci. 30 \(15\): 5437-50.](#)
8. Jean, Y.H. *et al.* (2009) Capnellene, a natural marine compound derived from soft coral, attenuates chronic constriction injury-induced neuropathic pain in rats [Br J Pharmacol. 158: 713-25.](#)
9. Morales-Garcia, J.A. *et al.* (2011) Phosphodiesterase 7 inhibition preserves dopaminergic neurons in cellular and rodent models of Parkinson disease. [PLoS One. 6\(2\):e17240.](#)
10. Spencer-Segal, J.L. *et al.* (2011) Distribution of Phosphorylated TrkB Receptor in the Mouse Hippocampal Formation Depends on Sex and Estrous Cycle Stage. [J Neurosci. 31: 6780-90.](#)
11. Chew, S.S. *et al.* (2011) Response of retinal Connexin43 to optic nerve injury. [Invest Ophthalmol Vis Sci. 52: 3620-9.](#)
12. Huh, S.H. *et al* (2011) Ethyl pyruvate rescues nigrostriatal dopaminergic neurons by regulating glial activation in a mouse model of Parkinson's disease. [J Immunol. 187: 960-9.](#)
13. Jeong, H.K. *et al.* (2010) Inflammatory responses are not sufficient to cause delayed neuronal death in ATP-induced acute brain injury. [PLoS One. 5: e13756.](#)
14. Liew, H.K. *et al.* (2012) Systemic administration of urocortin after intracerebral hemorrhage reduces neurological deficits and neuroinflammation in rats. [J Neuroinflammation. 9: 13.](#)
15. Szymdynger-Chodobska, J. *et al.* (2011) Multiple sites of vasopressin synthesis in the injured brain. [J Cereb Blood Flow Metab. 31: 47-51.](#)
16. Tchoukalova, Y.D. *et al.* (2012) *In vivo* adipogenesis in rats measured by cell kinetics in adipocytes and plastic-adherent stroma-vascular cells in response to high-fat diet and thiazolidinedione. [Diabetes. 61: 137-44.](#)
17. Xu, Q. *et al.* (2011) Spinal phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling cascades in inflammation-induced hyperalgesia. [J Neurosci. 31: 2113-24.](#)
18. Lovett-Barr, M.R. *et al.* (2012) Repetitive intermittent hypoxia induces respiratory and somatic motor recovery after chronic cervical spinal injury. [J Neurosci. 32 \(11\): 3591-600.](#)

19. Lavis, S. *et al.* (2012) Reactive astrocytes overexpress TSPO and are detected by TSPO positron emission tomography imaging. [J Neurosci. 32: 10809-18.](#)
20. Ortega, F.J. *et al.* (2012) Glibenclamide enhances neurogenesis and improves long-term functional recovery after transient focal cerebral ischemia. [J Cereb Blood Flow Metab. 33: 356-64.](#)
21. Zhao, H. *et al.* (2013) Brain 3-Mercaptopyruvate Sulfurtransferase (3MST): Cellular Localization and Downregulation after Acute Stroke. [PLoS One. 8\(6\):e67322.](#)
22. Huang S *et al.* (2015) Expression of Peroxiredoxin 1 After Traumatic Spinal Cord Injury in Rats. [Cell Mol Neurobiol. May 24. \[Epub ahead of print\]](#)
23. Wang, A.Y. *et al.* (2015) Searching for a rat model of chronic tympanic membrane perforation: Healing delayed by mitomycin C/dexamethasone but not paper implantation or iterative myringotomy. [Int J Pediatr Otorhinolaryngol. 79 \(8\): 1240-7.](#)
24. Hernangómez M *et al.* (2016) CD200R1 agonist attenuates glial activation, inflammatory reactions, and hypersensitivity immediately after its intrathecal application in a rat neuropathic pain model. [J Neuroinflammation. 13 \(1\): 43.](#)
25. Signarovitz, A.L. *et al.* (2012) Mucosal immunization with live attenuated *Francisella novicida* U112ΔiglB protects against pulmonary F. tularensis SCHU S4 in the Fischer 344 rat model. [PLoS One. 7: e47639.](#)
26. Liu, Z. *et al.* (2016) Leukocyte Infiltration Triggers Seizure Recurrence in a Rat Model of Temporal Lobe Epilepsy. [Inflammation. Apr 4. \[Epub ahead of print\].](#)
27. Szymdynger-Chodobska, J. *et al.* (2016) The Involvement of Pial Microvessels in Leukocyte Invasion after Mild Traumatic Brain Injury. [PLoS One. 11 \(12\): e0167677.](#)
28. Ejaz, S. *et al.* (2014) MRI and neuropathological validations of the involvement of air pollutants in cortical selective neuronal loss. [Environ Sci Pollut Res Int. 21 \(5\): 3351-62.](#)
29. Alizadeh, A. *et al.* (2017) Neuregulin-1 positively modulates glial response and improves neurological recovery following traumatic spinal cord injury. [Glia. Apr 29. \[Epub ahead of print\]](#)
30. Popiolek-Barczyk, K. *et al.* (2017) Biphalin, a Dimeric Enkephalin, Alleviates LPS-Induced Activation in Rat Primary Microglial Cultures in Opioid Receptor-Dependent and Receptor-Independent Manners. [Neural Plasticity. 2017: 1-19.](#)
31. Huang RY *et al.* (2017) Rapid and Delayed Effects of Pulsed Radiofrequency on Neuropathic Pain: Electrophysiological, Molecular, and Behavioral Evidence Supporting Long-Term Depression. [Pain Physician. 20 \(2\): E269-E283.](#)
32. Bourke, G. *et al.* (2017) Effects of early nerve repair on experimental brachial plexus injury in neonatal rats. [J Hand Surg \(Eur Vol\). 1753193417732696 \[Epub ahead of print\].](#)
33. Wang, Z.C. *et al.* (2017) Involvement of NF-κB and the CX3CR1 Signaling Network in Mechanical Allodynia Induced by Tetanic Sciatic Stimulation. [Neurosci Bull. Jun 13 \[Epub ahead of print\].](#)

Storage	<p>Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.</p> <p>DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL:RPE (MCA1210PE)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M300615:170106'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)