

Datasheet: MCA2801

Description:	MOUSE ANTI HUMAN MALT1
Specificity:	MALT1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	50
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			■	
Immunohistology - Frozen			■	
Immunohistology - Paraffin (1)	■			
ELISA			■	
Immunoprecipitation			■	
Western Blotting	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Recombinant protein corresponding to amino acids 701-808 located within the C-terminal of

MALT1.

**External Database
Links**

UniProt:

[Q9UDY8](#)

[Related reagents](#)

Entrez Gene:

[10892](#)

MALT1

[Related reagents](#)

Synonyms

MLT

Specificity

Mouse anti Human MALT1 antibody, clone 50 recognizes human mucosal-associated lymphoid tissue lymphoma translocation 1 (MALT1), also known as MALT lymphoma-associated translocation and paracaspase. MALT1 is an 824 amino acid ~93 kDa protein belonging to the [peptidase C14B family](#) containing a single [death domain](#) and two [Ig-like CD-type](#) domains. In normal lymphocytes, MALT1 plays an important role in antigen receptor-mediated lymphocyte activation. In T-cells, MALT1 is recruited by activated CARMA1, along with Bcl-10, to form a CARMA1-Bcl10-MALT1 (CBM) complex which is involved in the activation of NF-kappaB ([Yang et al. 2013](#)).

MALT1 was initially identified as a recurrent translocation t(11;18)(q21;q21) in mucosal-associated lymphomas ([Horsman et al. 1992](#)), which creates a functional fusion oncoprotein consisting of MALT1 and apoptosis inhibitor API2 ([Dierlam et al. 1999](#)). Overexpression of MALT1, or expression of the API2-MALT1 fusion protein, leads to the uncontrolled activation of NF-kappaB which is stimulus-independent. ([Zhou et al. 2005](#))

Mouse anti Human MALT1 recognizes both MALT1 and the MALT1-API2 fusion protein ([Murga Penas et al. 2007](#)).

References

1. Ye, H. et al. (2005) MALT lymphoma with t(14;18)(q32;q21)/IGH-MALT1 is characterized by strong cytoplasmic MALT1 and BCL10 expression. [J Pathol. 205 \(3\): 293-301.](#)
2. Coornaert, B. et al. (2008) T cell antigen receptor stimulation induces MALT1 paracaspase-mediated cleavage of the NF-kappaB inhibitor A20. [Nat Immunol. 9 \(3\): 263-71.](#)
3. Rebeaud, F. et al. (2008) The proteolytic activity of the paracaspase MALT1 is key in T cell activation. [Nat Immunol. 9 \(3\): 272-81.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:

10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)

[RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
 Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®549](#),
[DyLight®649](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
 Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
 Goat Anti Mouse IgG (STAR77...) [HRP](#)
 Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
 Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
 Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
 Goat Anti Mouse IgG (STAR70...) [FITC](#)
 Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
 Human Anti Mouse IgG1 (HCA036...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Recommended Useful Reagents

[HISTAR DETECTION SYSTEM \(STAR3000B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M315880:180503'

Printed on 05 May 2018