

Datasheet: MCA43PE

Description:	MOUSE ANTI RAT CD45:RPE
Specificity:	CD45
Other names:	LCA
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	OX-1
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 1 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1%	Bovine Serum Albumin	
	5%	Sucrose	
Immunogen	Rat thymocyte membrane glycoproteins.		

External Database Links

UniProt:

[P04157](#) [Related reagents](#)

Entrez Gene:

[24699](#) Ptpcr [Related reagents](#)

Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	Mouse anti Rat CD45 antibody, clone OX-1 recognizes CD45, also known as the leucocyte common antigen (LCA). The leucocyte common antigen consists of a family of heavily glycosylated membrane glycoproteins of molecular weight 180 – 240kDa. Antibodies recognising a common epitope on all of these isoforms are termed CD45, whilst those recognising only individual isoforms are termed CD45RA, CD45RO etc. OX-1 reacts with all forms of CD45 expressed by all haematopoietic cells, except erythrocytes. CD45 isoforms play complex roles in T-cell and B-cell antigen receptor signal transduction. This product is routinely tested in flow cytometry on rat splenocytes
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Sunderland, C.A. <i>et al.</i> (1979) Purification with monoclonal antibody of a predominant leukocyte-common antigen and glycoprotein from rat thymocytes. Eur J Immunol. 9 (2): 155-9. 2. Woollett, G.R. <i>et al.</i> (1985) Molecular and antigenic heterogeneity of the rat leukocyte-common antigen from thymocytes and T and B lymphocytes. Eur J Immunol. 15 (2): 168-73. 3. Martín, A. <i>et al.</i> (1995) Passive dual immunization against tumour necrosis factor-α (TNF-α) and IL-1 beta maximally ameliorates acute aminonucleoside nephrosis. Clin Exp Immunol. 99 (2): 283-8. 4. Sato, K. <i>et al.</i> (2001) Carbon monoxide generated by heme oxygenase-1 suppresses the rejection of mouse-to-rat cardiac transplants. J Immunol. 166 (6): 4185-94. 5. Murakami, K. <i>et al.</i> (2000) Regulation of mast cell signaling through high-affinity IgE receptor by CD45 protein tyrosine phosphatase. Int Immunol. 12 (2): 169-76. 6. Standring, R. <i>et al.</i> (1978) The predominant heavily glycosylated glycoproteins at the surface of rat lymphoid cells are differentiation antigens. Eur J Immunol. 8 (12): 832-9. 7. Giezeman-Smits, K.M. <i>et al.</i> (1999) The regulatory role of CD45 on rat NK cells in target cell lysis. J Immunol. 163 (1): 71-6. 8. Zilka, N. <i>et al.</i> (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. J Neuroimmunol. 209 (1-2): 16-25. 9. Schupp, N. <i>et al.</i> (2011) Mineralocorticoid receptor-mediated DNA damage in kidneys of DOCA-salt hypertensive rats. FASEB J. 25 (3): 968-78. 10. Ermert, L. <i>et al.</i> (2001) Comparison of different detection methods in quantitative microdensitometry. Am J Pathol. 158: 407-17. 11. Jeong, H.K. <i>et al.</i> (2010) Inflammatory responses are not sufficient to cause delayed neuronal death in ATP-induced acute brain injury. PLoS One. 5: e13756. 12. Leonardo, C.C. <i>et al.</i> (2009) Inhibition of gelatinase activity reduces neural injury in an ex vivo model of hypoxia-ischemia. Neuroscience. 160: 755-66. 13. Markusic, D.M. <i>et al.</i> (2010) Separating lentiviral vector injection and induction of gene expression in time, does not prevent an immune response to rTA in rats. PLoS One. 5: e9974. 14. Vaschetto, R. <i>et al.</i> (2010) Renal hypoperfusion and impaired endothelium-dependent vasodilation in an animal model of VILI: the role of the peroxynitrite-PARP pathway Crit Care. 14: R45. 15. Ladhoff, J. <i>et al.</i> (2010) Immune privilege of endothelial cells differentiated from endothelial progenitor cells. Cardiovasc Res. 88: 121-9. 16. Yao, Y. <i>et al.</i> (2016) Alendronate Attenuates Spinal Microglial Activation and Neuropathic Pain. J Pain. 17 (8): 889-903.

17. Wang, C. *et al.* (2015) Small activating RNA induces myogenic differentiation of rat adipose-derived stem cells by upregulating MyoD. [Int Braz J Urol. 41 \(4\): 764-72.](#)
18. Tanner, D.C. *et al.* (2015) cFLIP is critical for oligodendrocyte protection from inflammation. [Cell Death Differ. 22 \(9\): 1489-501.](#)
19. Runesson, E. *et al.* (2015) Nucleostemin- and Oct 3/4-positive stem/progenitor cells exhibit disparate anatomical and temporal expression during rat Achilles tendon healing. [BMC Musculoskelet Disord. 16: 212.](#)

Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA1209PE\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	--	------------------	--	---------------	--

'M300890:170106'

Printed on 05 May 2018