

Datasheet: MCA453GA

Description:	MOUSE ANTI RAT TCR ALPHA/BETA
Specificity:	TCR ALPHA/BETA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	R73
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/50 - 1/100
Immunohistology - Frozen	■			1/10 - 1/100
Immunohistology - Paraffin			■	
Immunohistology - Resin			■	
ELISA			■	
Immunoprecipitation	■			
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat
Species Cross Reactivity	Reacts with: Monkey, Cynomolgus monkey N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Immunogen	Rat T Blasts and Erythrocytes.
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the X63.Ag8.653 mouse myeloma cell line.
Specificity	Mouse anti Rat TCR alpha/beta antibody, clone R73 recognizes a constant determinant on the beta chain of the rat alpha/beta T cell receptor, expressed by 97% of peripheral rat T cells as defined by the OX-52 marker. R73 is mitogenic for unseparated spleen cells and for purified T cells. In the rat thymus, mature medullary cells express the R73 determinant at the same levels as peripheral T cells, whereas 94% of CD4 - CD8 - thymocytes are R73 negative. Mouse anti Rat TCR alpha/beta antibody, clone R73 is reported to stimulate adhesion between Thymic Dendritic Cells and Thymocytes (Colic et al. 2010).
Flow Cytometry	Use 10ul of the suggested working dilution to label 100ul of whole blood or 10 ⁶ cells in 100ul. Method sheets are available on request.
References	- Hünig, T. <i>et al.</i> (1989) A monoclonal antibody to a constant determinant of the rat T cell antigen receptor that induces T cell activation. Differential reactivity with subsets of immature and mature T lymphocytes. J Exp Med. 169 (1): 73-86. - Tomida, S. <i>et al.</i> (1994) Intercellular adhesion molecule-1 and leukocyte function-associated antigen-1 are involved in protection mediated by CD3+TCR alpha beta- T cells at the early stage after infection with <i>Listeria monocytogenes</i> in rats. Int Immunol. 6 (7): 955-61. - Colić, M. <i>et al.</i> (1996) Mechanisms involved in the binding of thymocytes to rat thymic dendritic cells. Dev Immunol. 5 (1): 37-51. - Kanellis, J. <i>et al.</i> (2010) JNK signalling in human and experimental renal ischaemia/reperfusion injury. Nephrol Dial Transplant. 25: 2898-908. - Nave, H. <i>et al.</i> (2008) Resistance of Janus kinase-2 dependent leptin signaling in natural killer (NK) cells: a novel mechanism of NK cell dysfunction in diet-induced obesity. Endocrinology. 149: 3370-8. - Tsuchida, M. <i>et al.</i> (1994) Identification of CD4- CD8- alpha beta T cells in the subarachnoid space of rats with experimental autoimmune encephalomyelitis. A possible route by which effector cells invade the lesions. Immunology. 81 (3): 420-7. - Matsumoto, Y. <i>et al.</i> (1994) Successful prevention and treatment of autoimmune encephalomyelitis by short-term administration of anti-T-cell receptor alpha beta antibody. Immunology. 81 (1): 1-7. - Pilipović, I. <i>et al.</i> (2010) Glucocorticoids, master modulators of the thymic catecholaminergic system? Braz J Med Biol Res. 43 (3): 279-84. - Milicevic, N.M. <i>et al.</i> (2005) T cells are required for the peripheral phase of B-cell maturation. Immunology. 116: 308-17. - Kenny, E. <i>et al.</i> (2000) Phenotypic analysis of peripheral CD4+ CD8+ T cells in the rat. Immunology. 101: 178-84. - Petrovic-Dergovic, D.M. <i>et al.</i> (2004) Somatostatin-14 alters the thymus size and relation among the thymocyte subpopulations in peripubertal rats. Neuropeptides. 38: 25-34. - Trinh, L. <i>et al.</i> (2008) The corneal endothelium in an endotoxin-induced uveitis model: correlation between in vivo confocal microscopy and immunohistochemistry. Mol Vis. 14: 1149-56. - Khalife, S. <i>et al.</i> (2016) Relationship Between <i>Pneumocystis carinii</i> Burden and the Degree of Host Immunosuppression in an Airborne Transmission Experimental Model. J Eukaryot Microbiol. 63 (3): 309-17. - Bat, E. <i>et al.</i> (2013) Physical properties and erosion behavior of poly(trimethylene carbonate-co-ε-caprolactone) networks. Macromol Biosci. 13 (5): 573-83. - Jörns, A. <i>et al.</i> (2015) TNF-α Antibody Therapy in Combination With the T-Cell-Specific Antibody Anti-TCR Reverses the Diabetic Metabolic State in the LEW.1AR1-iddm Rat. Diabetes. 64

(8): 2880-91.

16. Ahn, M. *et al.* (2015) Immunohistochemical study of Krüppel-like factor 4 in the spinal cords of rats with experimental autoimmune encephalomyelitis. [Acta Histochem. 117 \(6\): 521-7.](#)

17. Jörns A *et al.* (2014) Anti-TCR therapy combined with fingolimod for reversal of diabetic hyperglycemia by β cell regeneration in the LEW.1AR1-iddm rat model of type 1 diabetes. [J Mol Med \(Berl\). 92 \(7\): 743-55.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®549 , DyLight®649 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Human Anti Mouse IgG1 (HCA036...)	HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M315934:180503'

Printed on 05 May 2018