

Datasheet: MCA4644F

Description:	MOUSE ANTI HUMAN CD155:FITC
Specificity:	CD155
Other names:	POLIOVIRUS RECEPTOR
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	SKII.4
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
Immunogen	Human SK-N-S1 neuroblastoma cells.		
External Database Links	UniProt: P15151 Related reagents		

Entrez Gene:[5817](#) PVR [Related reagents](#)

Synonyms	PVS
Specificity	Mouse anti Human CD155 antibody, clone SKII.4 recognizes human CD155, otherwise known as poliovirus receptor (PVR). CD155 is a 397 amino acid type I transmembrane glycoprotein with a predicted molecular mass of ~45 kDa, a member of the nectin family, which acts as a mediator of cell-cell adhesion and cell migration, and is a receptor for poliovirus and cytomegalovirus. CD155 is a primate-restricted marker expressed by monocytes, macrophages, CD34+ thymocytes, endothelial and epithelial cells, and exists as both cell surface (alpha and delta) and secreted (beta and gamma) isoforms. CD155 promotes natural killer (NK) cell adhesion and NK effector functions, interacting with the NK receptors CD96 (Tactile) and CD226 (DNAM-1). CD155 forms a heterodimer with CD113 (nectin-3) which then binds to integrin alphaVbeta3. Mouse anti Human CD155 antibody, clone SKII.4 has been shown to inhibit NK cell-mediated lysis (Fuchs <i>et al.</i> 2004).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	1. Fuchs, A. <i>et al.</i> (2005) Paradoxical inhibition of human natural interferon-producing cells by the activating receptor NKp44. Blood. 106 (6): 2076-82. 2. Boles, K.S. <i>et al.</i> (2009) A novel molecular interaction for the adhesion of follicular CD4 T cells to follicular DC. Eur J Immunol. 39 (3): 695-703. 3. Cella, M. <i>et al.</i> (2010) Loss of DNAM-1 contributes to CD8+ T-cell exhaustion in chronic HIV-1 infection. Eur J Immunol. 40 (4): 949-54.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com 'M301331:170109'	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

Printed on 01 Aug 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)