

Datasheet: MCA4645FT

Description:	MOUSE ANTI HUMAN CD319:FITC
Specificity:	CD319
Other names:	CRACC
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	162
Isotype:	IgG2b
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
Immunogen	CD319 - HuIgG fusion protein.		
External Database Links	UniProt: Q9NQ25 Related reagents		

Entrez Gene:[57823](#) SLAMF7 [Related reagents](#)

Synonyms	CS1
Specificity	Mouse anti Human CD319 antibody, clone 162 recognizes human CD319, otherwise known as CRACC (CD2-like receptor-activating cytotoxic cells), a type I transmembrane protein and member of the CD2 receptor family, expressed by natural killer (NK) cells, cytotoxic lymphocytes and activated B cells. Unlike the CD2 family receptors 2B4 and NTB-A, which trigger NK cell-mediated cytotoxicity through the recruitment of the adaptor protein SAP (SLAM-associated protein); CD319 has been shown to activate cytotoxicity through a unique SAP-independent ERK-mediated signalling pathway, through association with, and subsequent phosphorylation by, the adaptor protein EAT-2.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	1. Bouchon, A. <i>et al.</i> (2001) Activation of NK cell-mediated cytotoxicity by a SAP-independent receptor of the CD2 family. J Immunol. 167 (10): 5517-21. 2. Tassi, I. & Colonna, M. (2005) The cytotoxicity receptor CRACC (CS-1) recruits EAT-2 and activates the PI3K and phospholipase Cgamma signaling pathways in human NK cells. J Immunol. 175 (12): 7996-8002. 3. Kawano, Y. <i>et al.</i> (2013) Hypoxia reduces CD138 expression and induces an immature and stem cell-like transcriptional program in myeloma cells. Int J Oncol. 43 (6): 1809-16. 4. Pojero, F. <i>et al.</i> (2016) Utility of CD54, CD229, and CD319 for the identification of plasma cells in patients with clonal plasma cell diseases. Cytometry B Clin Cytom. 90 (1): 91-100.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL:FITC \(MCA691F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

'M301533:170109'

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)