

Datasheet: MCA4646F

Description:	MOUSE ANTI HUMAN CD335:FITC
Specificity:	CD335
Other names:	NKp46
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	9E2
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
Immunogen	CD335-Fc fusion protein		
External Database Links	UniProt: O76036 Related reagents		

Entrez Gene:[9437](#) [NCR1](#) [Related reagents](#)

Synonyms	LY94
Specificity	Mouse anti Human CD335 antibody, clone 9E2 specifically recognizes human NK (natural killer) cell-specific marker CD335, otherwise known as NKp46, a 46kDa type I transmembrane glycoprotein, and the first identified member of the natural cytotoxicity receptor (NCR) family. CD335 is uniquely expressed by all resting and activated NK cells, and considered to be the major NK cell lysis receptor for autologous pathogen-infected and tumour target cells, during natural cytotoxicity responses. Ligands of CD335 include viral haemagglutinins (HAs), and heparan sulphate proteoglycans (HSPGs) on the surface of tumour cells. Mouse anti Human CD335 antibody, clone 9E2 mediates re-directed NK cell cytotoxicity (Yusa et al. 2002) and (Chen et al. 2007).
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	1. Chklovskaya, E. <i>et al.</i> (2004) Reconstitution of dendritic and natural killer-cell subsets after allogeneic stem cell transplantation: effects of endogenous flt3 ligand. Blood. 103 (10): 3860-8. 2. Yusa, S. <i>et al.</i> (2002) SHP-1- and phosphotyrosine-independent inhibitory signaling by a killer cell Ig-like receptor cytoplasmic domain in human NK cells. J Immunol. 168 (10): 5047-57. 3. Chen, Y. <i>et al.</i> (2007) Prostaglandin D2 suppresses human NK cell function via signaling through D prostanoid receptor. J Immunol. 179 (5): 2766-73. 4. Rutigliano, H.M. <i>et al.</i> (2016) Trophoblast Major Histocompatibility Complex Class I Expression Is Associated with Immune-Mediated Rejection of Bovine Fetuses Produced by Cloning. Biol Reprod. 95 (2): 39.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com "M301544:170109"	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)