

Datasheet: MCA461C

Description:	MOUSE ANTI HUMAN CD45RO:RPE-Cy5
Specificity:	CD45RO
Format:	RPE-CY5
Product Type:	Monoclonal Antibody
Clone:	UCHL1
Isotype:	IgG2a
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Chimpanzee, Marmoset N.B. Antibody reactivity and working conditions may vary between species		
Product Form	Purified IgG conjugated to RPE-Cy5 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE-Cy5 488nm laser	496	667
Preparation	Purified IgG prepared by affinity chromatography on Protein A from		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	0.2% Bovine Serum Albumin		
Immunogen	Human IL-2 dependent T cells		
External Database Links	UniProt: P08575 Related reagents Entrez Gene:		

Synonyms	CD45
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse P3/NS1/1-Ag4-1 myeloma cell line
Specificity	Mouse anti Human CD45RO monoclonal antibody, clone UCHL1 recognizes the low molecular weight isoform (180 kDa) of the leucocyte common antigen (LCA). The antigen is expressed by a functional subset of T cells with memory phenotype. In peripheral blood Mouse anti Human CD45RO, clone UCHL1 stains 40-80% of lymphocytes and all monocytes and granulocytes.
Flow Cytometry	Use 5ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	1. Smith, S.H. <i>et al.</i> (1986) Functional subsets of human helper-inducer cells defined by a new monoclonal antibody, UCHL1. Immunology 58: 63-70. 2. Norton, A.J. <i>et al.</i> (1986) Monoclonal antibody (UCHL1) that recognises normal and neoplastic T cells in routinely fixed tissues. J Clin Pathol. 39 (4): 399-405. 3. Beverley, P.C.L. <i>et al.</i> (1986) T-cell subsets and function. Progress in Immunology VI. Cinader, B., Miller, G.G., eds. Academic Press Orlando pp 941-948. 4. Beverley, P.C. (1987) Human T cell subsets. Immunol Lett. 14 (4): 263-7. 5. Terry, L.A. <i>et al.</i> (1987) Phenotypic heterogeneity of the CD4+ and CD8+ subsets. Leucocyte Typing III. McMichael, A.J., Beverley, P.C.L. <i>et al.</i> eds. University Press. pp 225-7. 6. Akbar, A.N. <i>et al.</i> (1988) Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T cells. J Immunol. 140 (7): 2171-8. 7. Terry, L.A. <i>et al.</i> (1988) The monoclonal antibody, UCHL1, recognizes a 180,000 MW component of the human leucocyte-common antigen, CD45. Immunology. 64 (2): 331-6. 8. Beverley, P.C.L. <i>et al.</i> (1988) Phenotypic diversity of the CD45 antigen and its relation to function. Immunology, Suppl. 1: 3-5. 9. Merckenschlager, M. <i>et al.</i> (1988) Limiting dilution analysis of proliferative responses in human lymphocyte populations defined by the monoclonal antibody UCHL1: implications for differential CD45 expression in T cell memory formation. Eur J Immunol. 18 (11): 1653-61. 10. Cavers, M. <i>et al.</i> (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. Clin Exp Immunol. 127: 60-5. 11. Hutnick, N.A. <i>et al.</i> (2010) Vaccination with Ad5 vectors expands Ad5-specific CD8 T cells without altering memory phenotype or functionality. PLoS One. 5: e14385. 12. Leigh, J.E. <i>et al.</i> (2006) Characterization of the immune status of CD8+ T cells in oral lesions of human immunodeficiency virus-infected persons with oropharyngeal Candidiasis. Clin Vaccine Immunol. 13: 678-83. 13. Nistala, K. <i>et al.</i> (2008) Interleukin-17-producing T cells are enriched in the joints of children with arthritis, but have a reciprocal relationship to regulatory T cell numbers. Arthritis Rheum. 58: 875-87. 14. Liu Y <i>et al.</i> (2015) Fractionation of human spermatogenic cells using STA-PUT gravity sedimentation and their miRNA profiling. Sci Rep. 5: 8084. 15. Wilson, C.L. <i>et al.</i> (2015) Ubiquitin C-terminal hydrolase 1: A novel functional marker for liver myofibroblasts and a therapeutic target in chronic liver disease. J Hepatol. 63 (6): 1421-8.
Storage	Store at +4°C.

DO NOT FREEZE

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	6 months from date of despatch.
Acknowledgements	Cy® and CyDye® are registered trademarks of GE Healthcare
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M299800:170104'

Printed on 05 May 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)