

Datasheet: MCA461GA

Description:	MOUSE ANTI HUMAN CD45RO
Specificity:	CD45RO
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	UCHL1
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/25 - 1/50
Immunohistology - Frozen	■			
Immunohistology - Paraffin	■			
ELISA			■	
Immunoprecipitation	■			
Western Blotting			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Chimpanzee, Marmoset N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml

Immunogen	Human IL-2 dependent T cells
External Database Links	UniProt: P08575 Related reagents Entrez Gene: 5788 PTPRC Related reagents
Synonyms	CD45
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse P3/NS1/1-Ag4-1 myeloma cell line
Specificity	Mouse anti Human CD45RO monoclonal antibody, clone UCHL1 recognizes the low molecular weight isoform (180 kDa) of the leucocyte common antigen (LCA). The antigen is expressed by a functional subset of T cells with memory phenotype. In peripheral blood Mouse anti Human CD45RO, clone UCHL1 stains 40-80% of lymphocytes and all monocytes and granulocytes.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
Immunohistology	This product does not require protein digestion pre-treatment of paraffin sections. This product does not require antigen retrieval using heat treatment prior to staining of paraffin sections.
Histology Positive Control Tissue	Tonsil
References	1. Smith, S.H. <i>et al.</i> (1986) Functional subsets of human helper-inducer cells defined by a new monoclonal antibody, UCHL1. Immunology 58: 63-70. 2. Norton, A.J. <i>et al.</i> (1986) Monoclonal antibody (UCHL1) that recognises normal and neoplastic T cells in routinely fixed tissues. J Clin Pathol. 39 (4): 399-405. 3. Beverley, P.C.L. <i>et al.</i> (1986) T-cell subsets and function. Progress in Immunology VI. Cinader, B., Miller, G.G., eds. Academic Press Orlando pp 941-948. 4. Beverley, P.C. (1987) Human T cell subsets. Immunol Lett. 14 (4): 263-7. 5. Terry, L.A. <i>et al.</i> (1987) Phenotypic heterogeneity of the CD4+ and CD8+ subsets. Leucocyte Typing III. McMichael, A.J., Beverley, P.C.L. <i>et al.</i> eds. University Press. pp 225-7. 6. Akbar, A.N. <i>et al.</i> (1988) Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T cells. J Immunol. 140 (7): 2171-8. 7. Terry, L.A. <i>et al.</i> (1988) The monoclonal antibody, UCHL1, recognizes a 180,000 MW component of the human leucocyte-common antigen, CD45. Immunology. 64 (2): 331-6. 8. Beverley, P.C.L. <i>et al.</i> (1988) Phenotypic diversity of the CD45 antigen and its relation to function. Immunology, Suppl. 1: 3-5. 9. Merckenschlager, M. <i>et al.</i> (1988) Limiting dilution analysis of proliferative responses in human lymphocyte populations defined by the monoclonal antibody UCHL1: implications for differential CD45 expression in T cell memory formation. Eur J Immunol. 18 (11): 1653-61. 10. Cavers, M. <i>et al.</i> (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. Clin Exp Immunol. 127: 60-5. 11. Hutnick, N.A. <i>et al.</i> (2010) Vaccination with Ad5 vectors expands Ad5-specific CD8 T cells without altering memory phenotype or functionality. PLoS One. 5: e14385. 12. Leigh, J.E. <i>et al.</i> (2006) Characterization of the immune status of CD8+ T cells in oral lesions

of human immunodeficiency virus-infected persons with oropharyngeal Candidiasis. [Clin Vaccine Immunol. 13: 678-83.](#)

13. Nistala, K. *et al.* (2008) Interleukin-17-producing T cells are enriched in the joints of children with arthritis, but have a reciprocal relationship to regulatory T cell numbers. [Arthritis Rheum. 58: 875-87.](#)

14. Liu Y *et al.* (2015) Fractionation of human spermatogenic cells using STA-PUT gravity sedimentation and their miRNA profiling. [Sci Rep. 5: 8084.](#)

15. Wilson, C.L. *et al.* (2015) Ubiquitin C-terminal hydrolase 1: A novel functional marker for liver myofibroblasts and a therapeutic target in chronic liver disease. [J Hepatol. 63 \(6\): 1421-8.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®549](#),
[DyLight®649](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Human Anti Mouse IgG2a (HCA037...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M315938:180503'

Printed on 05 May 2018