

Datasheet: MCA4635F

Description:	RAT ANTI MOUSE CD4:FITC
Specificity:	CD4
Other names:	L3T4 ANTIGEN, LY-4
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	GK1.5
Isotype:	IgG2b
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/100

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Murine CD4.		
External Database Links	UniProt: P06332 Related reagents		

Entrez Gene:

[12504](#) Cd4 [Related reagents](#)

Fusion Partners	Spleen cells from immunised Lewis rats were fused with cells of the SP2/0 myeloma cell line.
Specificity	Rat anti Mouse CD4 antibody, clone GK1.5 recognizes mouse CD4, a ~55 kDa protein also known as Ly-4 and L3T4. CD4 is a single chain transmembraneous glycoprotein which belongs to the immunoglobulin superfamily, and is primarily expressed on T helper cells, peripheral blood monocytes and tissue macrophages. CD4 is also expressed on a subpopulation of regulatory T cells (CD4⁺ CD25⁺), which play a key role in the maintenance of self tolerance. Rat anti Mouse CD4 antibody, clone GK1.5 has been reported to block CD4⁺ T-cell activation. It blocks class II MHC antigen-specific binding, thereby inhibiting functions such as class II MHC antigen-specific proliferation and the release of lymphokines. It may also be used for <i>in vivo</i> and <i>in vitro</i> cell depletion of CD4⁺ T-cells.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).
References	1. Dialynas, D.P. <i>et al.</i> (1983) Characterization of the murine T cell surface molecule, designated L3T4, identified by monoclonal antibody GK1.5: similarity of L3T4 to the human Leu-3/T4 molecule. J Immunol. 131 (5): 2445-51. 2. Wilde, D.B. <i>et al.</i> (1983) Evidence implicating L3T4 in class II MHC antigen reactivity; monoclonal antibody GK1.5 (anti-L3T4a) blocks class II MHC antigen-specific proliferation, release of lymphokines, and binding by cloned murine helper T lymphocyte lines. J Immunol. 131 (5): 2178-83. 3. Dialynas, D.P. <i>et al.</i> (1983) Characterization of the murine antigenic determinant, designated L3T4a, recognized by monoclonal antibody GK1.5: expression of L3T4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. Immunol Rev. 74: 29-56. 4. Zhou, Z. <i>et al.</i> (2011) Autoreactive marginal zone B cells enter the follicles and interact with CD4⁺ T cells in lupus-prone mice. BMC Immunol. 12: 7. 5. Näher, H. <i>et al.</i> (1985) Dynamics of T cells of L3T4 and Ly 2 phenotype within granulomas in murine listeriosis. Clin Exp Immunol. 60 (3): 559-64. 6. Ye, X. <i>et al.</i> (2000) Transient depletion of CD4 lymphocyte improves efficacy of repeated administration of recombinant adenovirus in the ornithine transcarbamylase deficient sparse fur mouse. Gene Ther. 7 (20): 1761-7. 7. Chu, N.R. <i>et al.</i> (2000) Immunotherapy of a human papillomavirus (HPV) type 16 E7-expressing tumour by administration of fusion protein comprising <i>Mycobacterium bovis</i> bacille Calmette-Guérin (BCG) hsp65 and HPV16 E7. Clin Exp Immunol. 121:216-25 8. Pletinckx, K. <i>et al.</i> (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3- IL-10⁺ regulatory T cells by engaging CD28 and CTLA-4. Eur J Immunol. 45 (2): 480-91. 9. Foy, S.P. <i>et al.</i> (2016) Poxvirus-Based Active Immunotherapy with PD-1 and LAG-3 Dual Immune Checkpoint Inhibition Overcomes Compensatory Immune Regulation, Yielding Complete Tumor Regression in Mice. PLoS One. 11 (2): e0150084. 10. Steinl, D.C. <i>et al.</i> (2016) Noninvasive Contrast-Enhanced Ultrasound Molecular Imaging Detects Myocardial Inflammatory Response in Autoimmune Myocarditis. Circ Cardiovasc Imaging. 9 (8): . 11. Olesen, M. N. <i>et al.</i> (2018) CD4 T cells react to local increase of α-synuclein in a pathology-associated variant-dependent manner and modify brain microglia in absence of brain pathology Heliyon. 4 (1): e00513.

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	--	------------------	--	---------------	--

'M301573:170109'

Printed on 01 Aug 2018