

Datasheet: MCA4635T

Description:	RAT ANTI MOUSE CD4
Specificity:	CD4
Other names:	L3T4 ANTIGEN, LY-4
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	GK1.5
Isotype:	IgG2b
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/10 - 1/200
Immunohistology - Frozen	■			
Immunohistology - Paraffin			■	
ELISA	■			
Immunoprecipitation	■			
Western Blotting	■			
Immunofluorescence	■			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Murine CD4.

**External Database
Links**

UniProt:

[P06332](#)

[Related reagents](#)

Entrez Gene:

[12504](#)

Cd4

[Related reagents](#)

Fusion Partners

Spleen cells from immunised Lewis rats were fused with cells of the SP2/0 myeloma cell line.

Specificity

Rat anti Mouse CD4 antibody, clone GK1.5 recognizes mouse CD4, a ~55 kDa protein also known as Ly-4 and L3T4. CD4 is a single chain transmembraneous glycoprotein which belongs to the immunoglobulin superfamily, and is primarily expressed on T helper cells, peripheral blood monocytes and tissue macrophages. CD4 is also expressed on a subpopulation of regulatory T cells (CD4⁺ CD25⁺), which play a key role in the maintenance of self tolerance.

Rat anti Mouse CD4 antibody, clone GK1.5 has been reported to block CD4⁺ T-cell activation. It blocks class II MHC antigen-specific binding, thereby inhibiting functions such as class II MHC antigen-specific proliferation and the release of lymphokines. It may also be used for *in vivo* and *in vitro* cell depletion of CD4⁺ T-cells.

Flow Cytometry

Use 10ul of the suggested working dilution to label 1x10⁶ cells in 100ul.

References

1. Dialynas, D.P. *et al.* (1983) Characterization of the murine T cell surface molecule, designated L3T4, identified by monoclonal antibody GK1.5: similarity of L3T4 to the human Leu-3/T4 molecule. [J Immunol. 131 \(5\): 2445-51.](#)
2. Wilde, D.B. *et al.* (1983) Evidence implicating L3T4 in class II MHC antigen reactivity; monoclonal antibody GK1.5 (anti-L3T4a) blocks class II MHC antigen-specific proliferation, release of lymphokines, and binding by cloned murine helper T lymphocyte lines. [J Immunol. 131 \(5\): 2178-83.](#)
3. Dialynas, D.P. *et al.* (1983) Characterization of the murine antigenic determinant, designated L3T4a, recognized by monoclonal antibody GK1.5: expression of L3T4a by functional T cell clones appears to correlate primarily with class II MHC antigen-reactivity. [Immunol Rev. 74: 29-56.](#)
4. Zhou, Z. *et al.* (2011) Autoreactive marginal zone B cells enter the follicles and interact with CD4⁺ T cells in lupus-prone mice. [BMC Immunol. 12: 7.](#)
5. Näher, H. *et al.* (1985) Dynamics of T cells of L3T4 and Ly 2 phenotype within granulomas in murine listeriosis. [Clin Exp Immunol. 60 \(3\): 559-64.](#)
6. Ye, X. *et al.* (2000) Transient depletion of CD4 lymphocyte improves efficacy of repeated administration of recombinant adenovirus in the ornithine transcarbamylase deficient sparse fur mouse. [Gene Ther. 7 \(20\): 1761-7.](#)
7. Chu, N.R. *et al.* (2000) Immunotherapy of a human papillomavirus (HPV) type 16 E7-expressing tumour by administration of fusion protein comprising *Mycobacterium bovis* bacille Calmette-Guérin (BCG) hsp65 and HPV16 E7. [Clin Exp Immunol. 121:216-25](#)
8. Pletinckx, K. *et al.* (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3⁺ IL-10⁺ regulatory T cells by engaging CD28 and CTLA-4. [Eur J Immunol. 45 \(2\): 480-91.](#)
9. Foy, S.P. *et al.* (2016) Poxvirus-Based Active Immunotherapy with PD-1 and LAG-3 Dual Immune Checkpoint Inhibition Overcomes Compensatory Immune Regulation, Yielding Complete Tumor Regression in Mice. [PLoS One. 11 \(2\): e0150084.](#)
10. Steinl, D.C. *et al.* (2016) Noninvasive Contrast-Enhanced Ultrasound Molecular Imaging Detects Myocardial Inflammatory Response in Autoimmune Myocarditis. [Circ Cardiovasc Imaging. 9 \(8\): .](#)
11. Olesen, M. N. *et al.* (2018) CD4 T cells react to local increase of α-synuclein in a pathology-associated variant-dependent manner and modify brain microglia in absence of brain pathology [Heliyon. 4 \(1\): e00513.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M319155:180719'

Printed on 01 Aug 2018