

Datasheet: MCA4701

Description:	HAMSTER ANTI MOUSE CD27
Specificity:	CD27
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	LG.3A10
Isotype:	IgG
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/20
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Species Cross Reactivity	Reacts with: Human, Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 0.5mg/ml
Immunogen	Armenian hamster fibroblast line ARHO12 transfected with mouse CD27 cDNA.

**External Database
Links**

UniProt:

[P41272](#) [Related reagents](#)
[P26842](#) [Related reagents](#)
[Q501W2](#) [Related reagents](#)

Entrez Gene:

[21940](#) Cd27 [Related reagents](#)
[939](#) CD27 [Related reagents](#)

Synonyms Tnfrsf7, TNFRSF7

Specificity **Hamster anti Mouse CD27 antibody, clone LG.3A10** recognizes mouse, rat and human CD27, a member of the tumour necrosis factor receptor superfamily. It is expressed in the thymus and spleen, but not in non-lymphoid tissues. CD27 acts as a ligand for CD70, which is expressed in T and B cells, and acts as a costimulatory signal in lymphocyte proliferation and immunoglobulin production.

CD27 lacks a death domain in its short cytoplasmic tail, but can induce apoptosis by binding with the apoptosis regulatory protein Siva (also known as CD27-binding protein).

Flow Cytometry Use 10ul of the suggested working dilution to label 1x10⁶ cells in 100ul.

References 1. Gravestein, L.A. *et al.* (1995) Novel mAbs reveal potent co-stimulatory activity of murine CD27. [Int Immunol. 7: 551-7.](#)

Storage Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

**Health And Safety
Information** Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight®549](#), [DyLight®649](#), [DyLight®800](#),
[FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

 Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 30 Apr 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)