

Datasheet: MCA4772EL

Description:	RAT ANTI MOUSE TREM-2:Low Endotoxin
Specificity:	TREM-2
Format:	Low Endotoxin
Product Type:	Monoclonal Antibody
Clone:	78.18
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			1/25 - 1/200
Immunohistology - Frozen			■	
Immunohistology - Paraffin			■	
ELISA			■	
Immunoprecipitation			■	
Western Blotting			■	
Immunofluorescence			■	
Functional Assays	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Endotoxin Level	<0.01EU/ug
Approx. Protein Concentrations	IgG concentration 1.0mg/ml

Immunogen	Mouse TREM-2 Fc fusion protein.
External Database Links	UniProt: Q99NH8 Related reagents Entrez Gene: 83433 Trem2 Related reagents
Synonyms	Trem2a, Trem2b, Trem2c
Fusion Partners	Lymph node cells from immunised Fisher rats were fused with cells of the rat YB-2/0 myeloma cell line.
Specificity	Rat anti Mouse Trem-2 antibody, clone 78.18 recognizes Triggering receptor expressed on myeloid cells 2 (TREM-2), an activating receptor of the Ig superfamily. TREM-2 forms a receptor signalling complex with DAP12 to trigger the activation of the immune response in macrophages and dendritic cells. It may be involved in chronic inflammation and may promote the production of constitutive rather than inflammatory chemokines and cytokines. TREM-2 may also have a functional role in osteoclast formation. Stimulation of TREM-2 with Rat anti Mouse Trem-2 antibody, clone 78.18 enhances multinucleated osteoclast formation <i>in vitro</i>. Rat anti Mouse Trem-2 antibody, clone 78.18 may also be used to block TREM-2 on pre-osteoclasts to inhibit the formation of multinuclear osteoclasts and to inhibit bone resorption by mature osteoclasts.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	1. Humphrey, M.B. <i>et al.</i> (2006) TREM2, a DAP12-associated receptor, regulates osteoclast differentiation and function. J Bone Miner Res. 21 (2): 237-45.
Storage	Store at -20°C only. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	18 months from date of despatch.
Acknowledgements	This product is for <i>in vitro</i> research use only. It is not to be used for commercial purposes. Use of this product to produce products for sale or for diagnostic, therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for commercial purposes, contact The Regents of the University of California.
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: 10162: https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®549 , DyLight®649 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP

Recommended Negative Controls

[RAT IgG1 NEGATIVE CONTROL:Low Endotoxin \(MCA1211EL\)](#)

North & South America	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

'M316033:180503'

Printed on 05 May 2018